



# Primate phylogenetic relationships and divergence dates inferred from complete mitochondrial genomes



Luca Pozzi<sup>a,b,c,\*</sup>, Jason A. Hodgson<sup>a,b,d</sup>, Andrew S. Burrell<sup>a</sup>, Kirstin N. Sterner<sup>e</sup>, Ryan L. Raaum<sup>b,f</sup>, Todd R. Disotell<sup>a,b</sup>

<sup>a</sup> Department of Anthropology, Center for the Study of Human Origins, New York University, New York, NY, United States

<sup>b</sup> New York Consortium in Evolutionary Primatology, United States

<sup>c</sup> Behavioral Ecology and Sociobiology Unit, German Primate Center, Göttingen, Germany

<sup>d</sup> Department of Life Sciences, Imperial College London, London, United Kingdom

<sup>e</sup> Department of Anthropology, University of Oregon, Eugene, OR, United States

<sup>f</sup> Department of Anthropology, Lehman College & The Graduate Center, City University of New York, Bronx, NY, United States

## ARTICLE INFO

### Article history:

Received 9 February 2013

Revised 17 February 2014

Accepted 19 February 2014

Available online 28 February 2014

### Keywords:

Mitogenomics  
Molecular clock  
Divergence date  
Fossil calibration  
Strepsirrhini  
Haplorhini

## ABSTRACT

The origins and the divergence times of the most basal lineages within primates have been difficult to resolve mainly due to the incomplete sampling of early fossil taxa. The main source of contention is related to the discordance between molecular and fossil estimates: while there are no crown primate fossils older than 56 Ma, most molecule-based estimates extend the origins of crown primates into the Cretaceous. Here we present a comprehensive mitogenomic study of primates. We assembled 87 mammalian mitochondrial genomes, including 62 primate species representing all the families of the order. We newly sequenced eleven mitochondrial genomes, including eight Old World monkeys and three strepsirrhines. Phylogenetic analyses support a strong topology, confirming the monophyly for all the major primate clades. In contrast to previous mitogenomic studies, the positions of tarsiers and colugos relative to strepsirrhines and anthropoids are well resolved. In order to improve our understanding of how fossil calibrations affect age estimates within primates, we explore the effect of seventeen fossil calibrations across primates and other mammalian groups and we select a subset of calibrations to date our mitogenomic tree. The divergence date estimates of the Strepsirrhine/Haplorhine split support an origin of crown primates in the Late Cretaceous, at around 74 Ma. This result supports a short-fuse model of primate origins, whereby relatively little time passed between the origin of the order and the diversification of its major clades. It also suggests that the early primate fossil record is likely poorly sampled.

© 2014 Elsevier Inc. All rights reserved.

## 1. Introduction

The order Primates, including 420 species and 73 genera (IUCN 2013.2 IUCN Red List of Threatened Species), is a major mammalian radiation with a rich diversity of morphological, behavioral, and ecological adaptations. Today the non-human members of this order are mainly restricted to tropical or subtropical regions of Asia, Central and South America, and Africa. The fossil record shows that primates were more widely distributed in the past,

including in North America and Europe (Covert, 2002). Despite the great interest of evolutionary biologists in this radiation, numerous issues about their origin and diversification remain unclear.

The traditional view of primate origins is that crown primates or euprimates (i.e., all descendants of the last common ancestor of the living species) originated and diversified soon after the Cretaceous–Paleogene (K–Pg) boundary (*explosive model*). According to this model, the abrupt extinction of dinosaurs and the radiation of flowering plants (angiosperms) opened a new set of ecological niches that placental mammals (Eutheria) and birds (Neaves) could exploit (Conroy, 1990; Rose, 2006). As a consequence, those groups of animals that remained fairly small and cryptic during the Cretaceous underwent a rapid adaptive radiation soon after the mass extinction 65 millions year ago (Rose, 2006; Wible et al., 2007; O’Leary et al., 2013).

\* Corresponding author at: Behavioral Ecology and Sociobiology Unit, German Primate Center, Kellnerweg 4, 37077 Göttingen, Germany.

E-mail addresses: [lpozzi@dpz.eu](mailto:lpozzi@dpz.eu) (L. Pozzi), [j.hodgson@imperial.ac.uk](mailto:j.hodgson@imperial.ac.uk) (J.A. Hodgson), [andrew.burrell@nyu.edu](mailto:andrew.burrell@nyu.edu) (A.S. Burrell), [ksterner@uoregon.edu](mailto:ksterner@uoregon.edu) (K.N. Sterner), [ryan.raaum@lehman.cuny.edu](mailto:ryan.raaum@lehman.cuny.edu) (R.L. Raaum), [todd.disotell@nyu.edu](mailto:todd.disotell@nyu.edu) (T.R. Disotell).

The main line of evidence supporting the explosive model is the absence of euprimate fossils prior to the K–Pg boundary. The earliest unambiguous fossil euprimates date from the beginning of the Eocene epoch and are no older than 56 Ma (Miller et al., 2005; Smith et al., 2006; Williams et al., 2010). The best-known representatives of these early euprimates are the two groups Omo- myoidea (e.g., *Teilhardina* and *Melanerimia*) and Adapoidea (e.g., *Cantius* and *Donrussellia*), possibly related to the living haplorhines and strepsirrhines, respectively (Gingerich, 1986; Martin, 1993; Ni et al., 2004; Miller et al., 2005; Smith et al., 2006; Rasmussen, 2007). The North African *Altiatlasius koulchii* (Sigé et al., 1990) is slightly earlier, dated to approximately 57 Ma in the late Paleocene, but is not unambiguously a euprimate (Rasmussen, 2007; Williams et al., 2010). Finally, the plesiadapiforms, which have in the past been included in the order Primates, are found in the fossil record between the latest Cretaceous and the early Eocene, with most specimens recovered from deposits dating to the Paleocene (56–66 Ma). However, in contemporary taxonomies, none of these fossils are included in the euprimates and are thought to represent either a stem-group that diverged prior to the origin of euprimates (Bloch and Silcox, 2001; Bloch and Boyer, 2002; Bloch et al., 2007; Silcox et al., 2007) or a non-primate mammalian group possibly related to primates or colugos (Martin, 1968; Cartmill, 1972; Wible and Covert, 1987; Beard and Houde, 1989; Beard, 1990, 1993; Kay and Thorington, 1990; Kay et al., 1992).

Analyses of molecular data have provided a quite different scenario for the origin and diversification of primates. Molecular studies have yielded divergence dates older than any known euprimate fossil, extending the origins of crown primates before the K–Pg boundary (63.7–95.0 Ma; Springer et al., 2003, 2012; Chatterjee et al., 2009; Fabre et al., 2009; Meredith et al., 2011; Perelman et al., 2011; Jameson et al., 2011; Wilkinson et al., 2011). This view of primate origins agrees with a number of recent molecular studies on mammal diversification suggesting a deep Cretaceous origin for most of the modern placental lineages (Madsen et al., 2001; Murphy et al., 2001; Springer et al., 2003; Bininda-Emonds et al., 2007; Meredith et al., 2011), despite the lack of fossils showing clear morphological traits of the crown group Placentalia (Wible et al., 2007; O’Leary et al., 2013). In contrast to the *explosive model*, these studies place the origin of eutherian orders in the Cretaceous, supporting either a *long-fuse model* (the origin of the orders extend deep into the Cretaceous, with the origin of the main intraordinal lineages near the K–Pg boundary) or a *short-fuse model* (both interordinal and intraordinal diversity originating deep in the Cretaceous) (Springer et al., 2003; Bininda-Emonds et al., 2007; Meredith et al., 2011).

While it is expected that molecules would overestimate and fossils underestimate divergence times (Steiper and Young, 2008), the early lineage origin dates inferred from molecular studies suggest that we are missing a substantial proportion of primate evolutionary history (~20–25 million years). A possible source of the discrepancy between paleontological and molecular models of primate origins lies in the calibration of molecular clocks. In order to convert molecular evolutionary distance estimates to chronological time, a set of well-dated fossils with well-supported phylogenetic placement is required. These calibration points are critical in molecular dating since they are the only source of information about absolute time (Steiper and Young, 2008; Ho and Phillips, 2009; Parham et al., 2012). Three main problems can affect the use of fossil information in estimating divergence times using molecular sequences: (1) the selection of inappropriate fossils; (2) the incompleteness of the fossil record in some areas of the tree; and (3) the phylogenetic misplacement of a fossil (e.g., placing a stem form in a crown group). Though these problems have long been recognized as critical in dating a molecular tree (Raaum et al., 2005; Rutschmann et al., 2007; Steiper and Young, 2008;

Ho and Phillips, 2009), the difficulty of accurately choosing proper calibration points is still underestimated (Pyrone, 2010; Dornburg et al., 2011; Parham et al., 2012). Many molecular studies suffer from a non-rigorous use of the paleontological record that may easily include considerable error in fossil calibration and provide misleading divergence estimates (Graur and Martin, 2004).

Here, we present one of the largest time-calibrated phylogenies of primates estimated using complete mitochondrial genome sequences. Mitogenomic analyses have been shown to be useful for phylogenetic reconstruction and divergence time estimation at different taxonomic levels and in different groups, including amphibians (Zardoya et al., 2003; Zhang et al., 2005; Zhang and Wake, 2009), birds (Pereira and Baker, 2006; Slack et al., 2007; Pacheco et al., 2011), fish (Inoue et al., 2003, 2010; Miya et al., 2003), and mammals (Arnason and Janke, 2002; Arnason et al., 2002, 2004, 2008). Mitochondrial genomes have traditionally been used for phylogenetic and phylogeographic analyses of animal taxa. The use of mitochondrial DNA (mtDNA) has several advantages including: the lack of recombination, a rapid coalescence time, relatively high substitution rates, high copy number, and haploidy. For these reasons, it has been suggested that mitochondrial gene trees are more likely to approximate the history of divergence among species than other loci (Moore, 1995). Today, multilocus phylogenies are very common (Jameson et al., 2011; Perelman et al., 2011; Springer et al., 2012), but the nuclear loci used in these studies are usually quite short (<1500 bp) compared to the large size of mitochondrial genomes (>16,000 bp). Mitochondrial genomes also show higher information content per base than nuclear DNA (Anderson et al., 1982; Cummings et al., 1995). Finally, the high number of copies of mitochondria in each cell makes mitogenomic data easy to obtain and sequence, especially in low quality samples such as museum and fossil specimens (Briggs et al., 2009; Krause et al., 2010; Rowe et al., 2011; Mason et al., 2011; Guschanski et al., 2013). Therefore, a good knowledge of the mitochondrial evolutionary history is not only valuable per se, but it can also provide an important comparative dataset in studies for which mtDNA is the only genetic marker available.

Within primates, mitochondrial genomes have been commonly used in phylogenetic studies at the family or genus level (Raaum et al., 2005; Sterner et al., 2006; Hodgson et al., 2009; Chan et al., 2010; Chiou et al., 2011; Guschanski et al., 2013; Zinner et al., 2013), but to date only two comprehensive mitogenomic studies including all main primate lineages have been performed (Matsui et al., 2009; Finstermeier et al., 2013). Here we assembled whole mitochondrial genome sequences from sixty-two primate species (including eleven new sequences) representing all living families within the order Primates. The goals of this study are (i) to infer the evolutionary relationships of primates using whole mitochondrial genome sequences, (ii) to investigate how different fossil calibrations selected within and outside primates affect estimates of times of divergence within the order, and (iii) to develop a suite of congruent fossil calibrations for use in future studies.

## 2. Materials and methods

### 2.1. Taxon sampling

We sequenced complete mitochondrial genomes for eleven species representing three strepsirrhines (*Daubentonia madagascarensis*, *Mirza coquereli*, and *Otolemur crassicaudatus*) and eight Old World monkeys (*Allenopithecus nigroviridis*, *Cercopithecus diana*, *C. lhoesti*, *C. mitis*, *Erythrocebus patas*, *Cercocebus torquatus*, *Lophocebus aterrimus*, and *Mandrillus sphinx*) (Table 1). Complete genome sequences from these specimens are deposited in GenBank

**Table 1**  
Sequences used in this study.

Scientific name	Common name	NCBI accession no	Reference
<b>Primates</b>			
Strepsirrhini			
Lorisiformes			
Lorisidae			
<i>Loris tardigradus</i>	Slender loris	NC_012763	Matsui et al. (2009)
<i>Nycticebus coucang</i>	Slow loris	NC_002765	Arnason et al. (2000)
<i>Perodicticus potto</i>	Potto	NC_012764	Matsui et al. (2009)
Galagidae			
<i>Galago senegalensis</i>	Senegal galago	NC_012761	Matsui et al. (2009)
<b><i>Otolemur crassicaudatus</i><sup>a</sup></b>	<b>Thick-tailed galago</b>	<b>KJ434961</b>	<b>This study</b>
Chiromyiformes			
Daubentoniidae			
<b><i>Daubentonia madagascariensis</i><sup>a</sup></b>	<b>Aye-aye</b>	<b>KJ158462</b>	<b>This study</b>
Lemuriformes			
Lemuridae			
<i>Eulemur mongoz</i>	Mongoose lemur	NC_010300	Arnason et al. (2008)
<i>Eulemur fulvus fulvus</i>	Brown lemur	NC_012766	Matsui et al. (2009)
<i>Eulemur fulvus mayottensis</i>	Brown lemur	NC_012769	Matsui et al. (2009)
<i>Eulemur macaco macaco</i>	Black lemur	NC_012771	Matsui et al. (2009)
<i>Lemur catta</i>	Ring-tailed lemur	NC_004025	Arnason et al. (2002)
<i>Varecia variegata</i>	Black and white ruffed lemur	NC_012773	Matsui et al. (2009)
Cheirogaleidae			
<b><i>Mirza coquereli</i><sup>a</sup></b>	<b>Coquereli's giant mouse lemur</b>	<b>KJ158463</b>	<b>This study</b>
Lepilemuridae			
<i>Lepilemur hubbardorum</i>	Hubbard's sportive lemur	NC_014453	Lei et al. (2010)
Indriidae			
<i>Propithecus coquereli</i>	Coquerel's sifaka	AB286049	Matsui et al. (2007)
Haplorhini			
Tarsiiformes			
Tarsiidae			
<i>Tarsius bancanus</i>	Horsfield's tarsier	NC_002811	Schmitz et al. (2002b)
<i>Tarsius syrichta</i>	Philippine tarsier	NC_012774	Matsui et al. (2009)
Anthropoidea			
Platyrrhini			
Cebidae			
<i>Aotus lemurinus</i>	Lemurine night monkey	FJ785421	Hodgson et al. (2009)
<i>Cebus albifrons</i>	White-fronted capuchin	AJ309866	Arnason et al. (2000)
<i>Saimiri sciureus</i>	Common squirrel monkey	FJ785425	Hodgson et al. (2009)
<i>Saguinus oedipus</i>	Cotton-top tamarin	FJ785424	Hodgson et al. (2009)
Pitheciidae			
<i>Callicebus donacophilus</i>	Bolivian titi	FJ785423	Hodgson et al. (2009)
Atelidae			
<i>Ateles belzebuth</i>	White-bellied spider monkey	FJ785422	Hodgson et al. (2009)
Catarrhini			
Cercopithecoidea			
Colobinae			
<i>Colobus guereza</i>	Guereza	AY863427	Raum et al. (2005)
<i>Nasalis larvatus</i>	Proboscis monkey	DQ355298	Sterner et al. (2006)
<i>Ptilocolobus badius</i>	Western red colobus	DQ355301	Sterner et al. (2006)
<i>Presbytis melalophos</i>	Mitred leaf monkey	DQ355299	Sterner et al. (2006)
<i>Pygathrix nemaeus</i>	Douc langur	DQ355302	Sterner et al. (2006)
<i>Rhinopithecus roxellana</i>	Golden snub-nosed monkey	DQ355300	Sterner et al. (2006)
<i>Semnopithecus entellus</i>	Hanuman langur	DQ355297	Sterner et al. (2006)
<i>Trachypithecus obscurus</i>	Dusky leaf monkey	NC_006900	Raum et al. (2005)
Cercopithecinae			
<b><i>Allenopithecus nigroviridis</i><sup>a</sup></b>	<b>Allen's swamp monkey</b>	<b>KJ434962</b>	<b>This study</b>
<b><i>Cercocebus torquatus</i><sup>a</sup></b>	<b>Collared mangabey</b>	<b>KJ434959</b>	<b>This study</b>
<b><i>Cercopithecus diana</i><sup>a</sup></b>	<b>Diana monkey</b>	<b>KJ434958</b>	<b>This study</b>
<b><i>Cercopithecus lhoesti</i><sup>a</sup></b>	<b>L'Hoest's monkey</b>	<b>KJ434957</b>	<b>This study</b>
<b><i>Cercopithecus mitis</i><sup>a</sup></b>	<b>Blue monkey</b>	<b>KJ434956</b>	<b>This study</b>
<i>Chlorocebus aethiops</i>	Grivet	NC_007009	Raum et al. (2005)
<i>Chlorocebus pygerythrus</i>	Vervet monkey	NC_009747	Wertheim and Worobey (2007)
<i>Chlorocebus sabaues</i>	Green monkey	EF597503	Wertheim and Worobey (2007)
<i>Chlorocebus tantalus</i>	Tantalus monkey	NC_009748	Wertheim and Worobey (2007)
<b><i>Erythrocebus patas</i><sup>a</sup></b>	<b>Patas monkey</b>	<b>KJ434955</b>	<b>This study</b>
<b><i>Lophocebus aterrimus</i><sup>a</sup></b>	<b>Black crested mangabey</b>	<b>KJ434960</b>	<b>This study</b>
<i>Macaca sylvanus</i>	Barbary ape	NC_002764	Arnason et al. (2000)
<i>Macaca mulatta</i>	Rhesus monkey	NC_005943	Gokey et al. (2004)
<i>Macaca fascicularis</i>	Crab-eating macaque	NC_012670	Yi et al. (unpub.)
<i>Macaca thibetana</i>	Pere David's macaque	NC_011519	Li et al. (2009)
<b><i>Mandrillus sphinx</i><sup>a</sup></b>	<b>Mandrill</b>	<b>KJ434963</b>	<b>This study</b>
<i>Papio hamadryas</i>	Hamadryas baboon	NC_001992	Arnason et al. (1998)
<i>Theropithecus gelada</i>	Gelada	FJ785426	Hodgson et al. (2009)

(continued on next page)

Table 1 (continued)

Scientific name	Common name	NCBI accession no	Reference
<b>Hominoidea</b>			
<i>Gorilla gorilla</i>	Western gorilla	NC_001645	Horai et al. (1995)
<i>Homo neanderthalensis</i>	Neanderthal	AM948965	Green et al. (2008)
<i>Homo sapiens</i>	Human	EF061150	Friedlaender et al. (2007)
<i>Homo sp.</i> (Denisovan)	Denisova hominin	FR695060	Krause et al. (2010)
<i>Hylobates agilis</i>	Agile gibbon	NC_014042	Matsudaira and Ishida (2010)
<i>Hylobates lar</i>	White-handed gibbon	NC_002082	Arnason et al. (1996)
<i>Hylobates pileatus</i>	Pileated gibbon	NC_014045	Matsudaira and Ishida (2010)
<i>Nomascus siki</i>	Southern white-cheeked gibbon	NC_014051	Matsudaira and Ishida (2010)
<i>Pan paniscus</i>	Pygmy chimpanzee	D38116	Horai et al. (1995)
<i>Pan troglodytes</i>	Common chimpanzee	EU095335	Flynn and Carr (2007)
<i>Pongo abelii</i>	Sumatran orangutan	X97707	Xu and Arnason (1996)
<i>Pongo pygmaeus</i>	Bornean orangutan	NC_001646	Horai et al. (1995)
<i>Symphalangus syndactylus</i>	Siamang	NC_014047	Matsudaira and Ishida (2010)
<b>Non-primate mammals</b>			
<b>Euarchontoglires</b>			
<b>Dermoptera</b>			
<i>Galeopterus variegatus</i>	Sunda colugo	NC_004031	Arnason et al. (2002)
<i>Galeopterus variegatus</i>	Sunda colugo	AF460846	Schmitz et al. (2002a)
<b>Rodentia</b>			
<i>Cavia porcellus</i>	Guinea pig	AJ222767	D'Erchia et al. (1996)
<i>Cricetulus griseus</i>	Chinese hamster	NC_007936	Partridge et al. (2007)
<i>Glis glis</i>	Fat dormouse	NC_001892	Reyes et al. (1998)
<i>Mus musculus</i>	Mouse	NC_005089	Bayona-Bafaluy (2003)
<i>Rattus norvegicus</i>	Brown rat	AJ428514	Nilsson et al. (2003)
<i>Sciurus vulgaris</i>	Squirrel	AJ238588	Reyes et al. (2000)
<b>Lagomorpha</b>			
<i>Lepus europaeus</i>	European hare	AJ421471	Arnason et al. (2002)
<i>Ochotona collaris</i>	Collared pika	NC_003033	Lin et al. (2002)
<i>Oryctolagus cuniculus</i>	Rabbit	NC_001913	Gissi et al. (1998)
<b>Laurasiatheria</b>			
<b>Carnivora</b>			
<i>Canis familiaris</i>	Dog	NC_002008	Kim et al. (1998)
<i>Felis catus</i>	Domestic cat	U20753	Lopez et al. (1996)
<b>Perissodactyla</b>			
<i>Equus caballus</i>	Horse	NC_001640	Xu and Arnason (1994)
<i>Rhinoceros unicornis</i>	Greater Indian rhinoceros	NC_001779	Xu et al. (1996)
<b>Cetartiodactyla</b>			
<i>Balaenoptera musculus</i>	Blue whale	NC_001601	Arnason and Gullberg (1993)
<i>Balaenoptera physalus</i>	Finback whale	X61145	Arnason et al. (1991)
<i>Bos taurus</i>	Cow	NC_006853	Chung and Ha (unpub).
<i>Hippopotamus amphibius</i>	Hippopotamus	AJ010957	Ursing and Arnason (1998b)
<i>Lama pacos</i>	Alpaca	NC_002504	Ursing et al. (2000)
<i>Ovis aries</i>	Sheep	AF010406	Hiendleder and Mainz (1998)
<i>Physeter catodon</i>	Sperm whale	NC_002503	Arnason et al. (2000)
<i>Sus scrofa</i>	Pig	AJ002189	Ursing and Arnason (1998a)
<b>Afrotheria</b>			
<b>Proboscidea</b>			
<i>Elephas maximus</i>	Asiatic elephant	NC_005129	Rogaev et al. (2006)
<i>Loxodonta africana</i>	African elephant	NC_000934	Hauf et al. (2000)

<sup>a</sup> In bold new sequences generated for this study.

(<http://www.ncbi.nlm.nih.gov/>) under accession numbers KJ158462–KJ158463 and KJ434955–KJ434963 (Table 1).

These genomes were added to a dataset including mitochondrial genomes of 51 other primates, in total representing all major taxonomic lineages within the order, as well as mitogenomes of 23 other species representing major lineages within Boreoeutheria. Rooting was done using two Afrotheria species, *Elephas maximus* and *Loxodonta africana*, commonly accepted as external to Boreoeutheria in mammal phylogenies (Murphy et al., 2001; Arnason and Janke, 2002; Springer et al., 2003; Bininda-Emonds et al., 2007; Arnason et al., 2008; Meredith et al., 2011; Song et al., 2012). Overall 87 mammalian genomes were included in the phylogenetic analyses.

## 2.2. Sequencing

DNA was extracted and isolated from either blood or tissue samples using the QIAamp DNA Micro Kit (Qiagen, Inc.) following

the protocol provided by the manufacturer. A list of the samples with location, sample type, and GenBank accession number is reported in Supplementary Table A1. All the samples are currently stored at the Molecular Anthropology lab at New York University. Mitochondrial genomes were sequenced by long-template PCR, which minimizes the chance of amplifying mitochondrial pseudogenes in the nuclear genome (*numts*) (Thalmann et al., 2004; Raaum et al., 2005). We employed the same approach previously used in Raaum et al. (2005), Sterner et al. (2006), Hodgson et al. (2009), and Chiou et al. (2011). Briefly, independent sets of amplification primers that amplify the entire mtDNA genome in two or more overlapping long fragments were designed for each species. PCR reactions were performed employing a long-range PCR kit (Expand Long Template PCR System, Roche). The sequencing products were analyzed on a 3730 DNA Analysis System (Applied Biosystems, Foster City, CA). Sequences were then assembled and edited using Sequencher 4.1 (Gene Codes, Corp., Ann Arbor, MI). In addition to the eleven newly sequenced genomes, fifteen of the primate

genomes used in this study were previously sequenced in the NYU Molecular Anthropology lab applying the technique described above (see Raalum et al., 2005; Sterner et al., 2006; Hodgson et al., 2009). All the remaining genomes (61) obtained from GenBank were analyzed to minimize the presence of nuclear copies of mitochondrial genes (*numts*) in the dataset. First, we translated all of the protein-coding mitochondrial genes into amino acids to check for the presence of premature stop codons and frame shifting insertions or deletions. Then, for each protein-coding gene and RNA sequence independently, we constructed a neighbor-joining tree to identify any sequence that displayed unusual phylogenetic placement.

### 2.3. Alignment

Heavy-strand protein-coding genes and RNA sequences (2 rRNAs and 22 tRNAs) were automatically extracted from complete genome sequences according to GenBank annotations using a Perl script written by JAH for this purpose. The D-loop was excluded from the dataset because of alignment difficulties due to its high variability. Also the ND6 gene was excluded from the final alignment because it is encoded on the mitochondrial L-strand which has a different nucleotide composition from the H-strand, and has been shown to have poor phylogenetic signal (Gissi et al., 2000; Miya and Nishida, 2000). Protein-coding genes were then aligned based on their corresponding amino acid translations using the software TranslatorX (Abascal et al., 2010). Since homology is best identified at the amino acid level due to the evolution of coding DNA as triplets of nucleotides and the redundancy of the genetic code, this software translates the DNA sequences into amino acids, aligns the amino acid sequences, and then back-translates the alignment to the nucleotide sequences. We allowed the software to automatically identify the most likely reading frame (minimizing the number of stop codons) and we used MUSCLE (Edgar, 2004) to perform the protein alignment. RNA sequences (rRNAs and tRNAs) were aligned with reference to their secondary structure. Due to their stem-and-loop structure, the identification of homologous regions is best identified when the secondary structure of the RNA sequences is taken into account. This approach has been advocated to improve not only the alignment itself but also the phylogenetic tree reconstruction (Kjer and Honeycutt, 2007; Stocsits et al., 2009; Letsch et al., 2010). We employed two different structure informed alignment methods for RNA sequences: RNAsalsa (Stocsits et al., 2009) and RAF (Do et al., 2006, 2008). RNAsalsa, specifically designed to align ribosomal RNAs, uses preexisting knowledge of RNA structure to simultaneously predict secondary structure in the RNA sequences and to align them based on the secondary structure information. We used secondary structures of *Bos taurus* as reported in the literature (Springer and Douzery, 1996; Burk and Douzery, 2002; Stocsits et al., 2009) to align the remaining 86 sequences in our dataset. RAF (RNA Alignment and Folding) uses an algorithm for simultaneous alignment and consensus folding of unaligned RNA sequences. In contrast to RNAsalsa, RAF does not require preexisting information about secondary structure but identifies sets of likely pairing and alignment candidates for each nucleotide to effectively obtain a simultaneous pairwise alignment and folding (Do et al., 2006, 2008) and then compares the results of the two candidate alignments to identify possible discrepancies.

The resulting alignments (12 protein coding genes, 12S and 16S rRNAs, and 22 tRNAs) were concatenated after removing problematic regions using Gblocks 0.91 (Talavera and Castresana, 2007) under a relaxed approach. Gblocks removes all poorly aligned regions in a dataset and it has been shown to be particularly effective in phylogenetic studies including very divergent sequences (Castresana, 2000; Talavera and Castresana, 2007). Gblocks selects

blocks to be deleted following a reproducible set of requirements, including the lack of large segments of contiguous non-conserved positions, lack of gap positions and high conservation of flanking positions. Gblocks was run with the options “Minimum Length Of A Block” = 10 (5 for RNAs), and “Allowed Gap Positions” = “With Half”.

A total of 11,022 bp from the 12 protein-coding genes, 1313 bp from 22 tRNA genes, and 1708 bp from two rRNA genes were unambiguously aligned. The individual alignments were then concatenated using the software SequenceMatrix v1.7.6 (Vaidya et al., 2011) to create a master alignment of 14,043 bp total, equivalent to approximately 83% of the mitochondrial genome.

### 2.4. Data partitions

For both Bayesian and maximum likelihood (ML) analyses, we evaluated six different partition schemes of the mitochondrial genome: (1) no partition (MT0); (2) four partitions (protein-coding genes, 12S rRNA, 16rRNAs, and 22 tRNAs) (MT4); (3) five partitions, identical to the previous one but with the protein-coding genes separated in two different partitions (i) codon position 1 and 2 and (ii) codon position 3 (MT5); (4) six partitions, consisting of three partitions for the two rRNAs (12S and 16S) and the 22 tRNAs, and three partitions for the three codon positions of the protein-coding genes (MT6); (5) fifteen partitions, where the 12 protein-coding genes were treated as independent partitions (MT15); and (6) 39 partitions, where the different codon positions of each protein-coding gene were treated as separate partition, plus the three RNA partitions (MT39). The nucleotide substitution model for each partition was selected among 24 models using the Akaike Information Criterion (AIC) (Posada and Buckley, 2004) as implemented in MrModeltest v2.3 (Nylander, 2004). For each of the data partitions, we evaluated the variable sites and the parsimony-informative sites across all taxa and within the ingroup (primates) only (Table 2).

### 2.5. Phylogenetic analysis

#### 2.5.1. Maximum likelihood (ML)

ML analyses were run using a Randomized Accelerated Maximum Likelihood in RAxML version 7.2.6 (Stamatakis et al., 2005; Stamatakis, 2006). For each partition scheme, we ran 50 independent ML inferences (using 50 distinct randomized MP trees) with a GTRGAMMA model to estimate the best topology. We then performed a rapid (-f a -x option) with 1000 replications to assess support on different nodes (Stamatakis, 2006; Stamatakis et al., 2008). Maximum-likelihood bootstrap proportions (MLBS)  $\geq 70\%$  were considered strong support (Hillis and Bull, 1993; Wilcox et al., 2002).

#### 2.5.2. Bayesian analyses (MB)

Bayesian phylogenetic analyses were conducted with MrBayes 3.2.1 (Huelsenbeck and Ronquist, 2001; Ronquist and Huelsenbeck, 2003) using the Metropolis coupled Markov Chain Monte Carlo (Metropolis coupled MCMC or MC<sup>3</sup>) algorithm. The most appropriate model of nucleotide evolution was estimated independently for each partition using the AIC test as implemented in MrModelTest 2.3 (Nylander, 2004) as reported in Table 2. Although the model selected in most partitions was GTR + G + I, we also run our analyses using a simplified model GTR + G. This was done because Bayesian analyses may run into problems of convergence of the MCMC when one tries to accommodate among-site substitution rate variation using a combination of proportion of invariant sites model (+I) and a gamma distributed rates model (+G). This conflict may induce two regions of high posterior probability creating clear problems of convergence in the Bayesian analysis

**Table 2**  
Sequence characteristics and models chosen for each data partition.

Partition	No. sites	All taxa		Only primates		Model
		VS	PIS	VS	PIS	
All genes	14,043	9080	8244	8396	7532	GTR + I + G
12S rRNA	691	349	308	286	244	GTR + I + G
16S rRNA	1017	489	389	419	313	GTR + I + G
All tRNAs	1313	777	643	652	514	GTR + I + G
All protein-coding genes	11,022	7465	6904	7039	6461	GTR + I + G
pos1	3674	2277	2033	2092	1827	GTR + I + G
pos2	3674	1568	1270	1341	1075	GTR + I + G
pos1 + 2	7348	3845	3303	3433	2902	GTR + I + G
pos3	3674	3620	3601	3606	3559	GTR + I + G
Individual protein-coding genes						
<i>ATP6</i>	681	499	465	477	434	GTR + I + G
pos1	227	163	151	156	134	GTR + I + G
pos2	227	112	92	98	80	GTR + I + G
pos3	227	224	222	223	220	GTR + I + G
<i>ATP8</i>	225	198	190	193	176	GTR + G
pos1	75	64	59	62	55	GTR + G
pos2	75	64	62	61	54	GTR + G
pos3	75	70	69	70	67	GTR + G
<i>COI</i>	1569	807	727	752	694	GTR + I + G
pos1	523	198	153	164	135	GTR + I + G
pos2	523	96	62	76	50	GTR + I + G
pos3	523	513	512	512	509	GTR + I + G
<i>COII</i>	696	443	408	417	385	GTR + I + G
pos1	232	134	121	124	111	GTR + I + G
pos2	232	81	60	66	49	HKY + I + G
pos3	232	228	227	227	225	GTR + I + G
<i>COIII</i>	804	463	426	443	406	GTR + I + G
pos1	268	131	110	122	101	GTR + I + G
pos2	268	72	58	62	50	GTR + I + G
pos3	268	260	258	259	255	GTR + I + G
<i>CYTB</i>	1143	747	961	714	648	GTR + I + G
pos1	381	221	200	209	182	GTR + I + G
pos2	381	147	114	126	102	HKY + I + G
pos3	381	379	377	379	364	GTR + I + G
<i>NADH1</i>	963	640	599	601	556	GTR + I + G
pos1	321	203	180	176	160	GTR + I + G
pos2	321	118	101	106	81	GTR + I + G
pos3	321	319	318	319	315	GTR + G
<i>NADH2</i>	1050	848	478	788	729	GTR + I + G
pos1	350	274	252	258	230	GTR + I + G
pos2	350	226	192	185	159	GTR + I + G
pos3	350	348	34	345	340	GTR + G
<i>NADH3</i>	360	248	231	234	219	GTR + I + G
pos1	120	78	71	69	64	GTR + I + G
pos2	120	54	45	50	42	GTR + I + G
pos3	120	116	115	115	113	GTR + G
<i>NADH4</i>	1386	1002	911	951	864	GTR + I + G
pos1	462	319	287	297	258	GTR + I + G
pos2	462	225	179	197	157	GTR + I + G
pos3	462	458	445	457	449	GTR + I + G
<i>NADH4L</i>	297	222	205	204	186	GTR + I + G
pos1	99	74	68	68	60	GTR + I + G
pos2	99	51	40	40	30	GTR + G
pos3	99	97	97	96	96	GTR + I + G
<i>NADH5</i>	1848	1348	1253	1265	1154	GTR + I + G
pos1	616	418	381	387	337	GTR + I + G
pos2	616	322	265	274	221	GTR + I + G
pos3	616	608	607	604	596	GTR + I + G

Note: VS = variable sites; PIS = parsimony-informative sites.

(Brian Moore, pers. comm.; Stamatakis, 2006). Four independent runs, each including four incrementally heated chains, were run for at least 20 million generations. Different partition schemes required different number of generations to properly reach convergence as reported in Table 3. The sample frequency was adjusted according to the number of generations (Table 3). We assessed convergence visually using Tracer v.1.5 (Rambaut and Drummond, 2007) to plot the likelihood versus generation number and estimate the effective sample size (ESS > 200) of all parameters and to compare the performance of the four independent analyses. In

addition, we used AWTY (Are We There Yet?, Nylander et al., 2008) to plot pairwise split frequencies for the four independent MCMC runs and to check the posterior probabilities of clades for non-overlapping samples of trees in the sample using the compare and slide commands, respectively. After checking for convergence we summarized the posterior distribution of trees removing the first 10% of generations as burn-in. Posterior probability (PP) support values higher than 0.95 were considered strong support for individual clades (e.g., Alfaro, 2003; Erixon et al., 2003; Huelsenbeck and Rannala, 2004).

**Table 3**  
Summary of model partition comparison (maximum likelihood and Bayesian analyses).

Model	Maximum likelihood	Bayesian				
		Run	Generations	Burn-in	Marginal likelihood	SD
MT0	−421145.742	1	20,000,000	2,000,000	−417245.466	0.239
		2	20,000,000	2,000,000	−417245.834	0.287
MT4	−420115.955	1	20,000,000	2,000,000	−415805.630	0.329
		2	20,000,000	2,000,000	−415804.224	0.280
MT5	−409906.041	1	20,000,000	2,000,000	−409740.282	0.295
		2	20,000,000	2,000,000	−409740.561	0.329
MT6	−408934.350	1	20,000,000	2,000,000	−408453.105	0.300
		2	20,000,000	2,000,000	−408453.716	0.297
MT15	−419192.485	1	30,000,000	3,000,000	−414803.673	0.311
		2	30,000,000	3,000,000	−414814.001	0.381
MT39	−407068.508	1	40,000,000	4,000,000	−402921.612	0.318
		2	40,000,000	4,000,000	−402921.590	0.364

## 2.6. Partition models comparison

In order to evaluate the relative support for competing partition models, we compared the marginal likelihood scores for both maximum likelihood and Bayesian analyses (Table 3). We also used Bayes factors (Kass and Raftery, 1995) to select the partition model better supported by Bayesian analyses. We calculated  $\ln$  (Bayes factor) as the difference in estimated marginal likelihoods for pairs of models, as described in (Nylander et al., 2004). For Bayesian analyses, marginal likelihoods were estimated from the harmonic means using Tracer v1.5 (Rambaut and Drummond, 2007) by employing the weighted likelihood bootstrap estimator with standard error estimated using 1000 bootstrap pseudoreplicates (Newton, 1994; Suchard et al., 2001). The results for the partition model comparison reported here are based on combining the post-burn-in data from the two runs that achieved the greatest marginal likelihoods, as summarized in Table 3. The similarity of tree topologies resulting from the analyses using different partitions was assessed using the program TOPD-FMITS version 3.3 (Puigbò et al., 2007). Unrooted maximum likelihood and Bayesian trees (without branch lengths) for each of the six partitioning schemes were compared. The TOPD-FMITS analysis was performed under default conditions, with 100 simulated trees as a null model. Analyses were run using the *nodal*, the *split*, and the *disagree* options. The *nodal method* compares the number of nodes that separate each taxon from the other taxa in the tree and calculates pairwise distance matrices from two input trees using only those taxa that are common to both trees (Puigbò et al., 2007). The *split* option uses a crossover method that takes into account the minimum number of operations required to convert one tree into the other, following the algorithm proposed by Robinson and Foulds (1981). Finally, the *disagree* option compares two trees and returns the taxa that make the disagreement between those trees. It first removes one taxon and calculates the gain (reduction in the split distance) between two trees. The taxon that produces most gain is removed for the following iterations. Then this procedure is repeated until the split distance is zero (Puigbò et al., 2007).

## 2.7. Test for compositional biases

Research over the last decade has shown that phylogenetic inference can be affected by systematic error, such as non-phylogenetic signals that can drive phylogenetic results (Phillips et al., 2004; Phillips, 2009). Even complex models of nucleotide substitution may not be able to capture the substitution process well enough to reliably provide an accurate phylogenetic reconstruction (Phillips et al., 2004; Phillips, 2009). This problem might be even more relevant in large concatenated datasets in which “long sequences exacerbate the potential for biases (systematic error) to be positively misleading” (Phillips et al., 2004: p. 1455).

The encoding of nucleotides as purines and pyrimidines (RY-coding) has been suggested as an effective approach to assess the effects of compositional biases and non-phylogenetic signal (multiple substitutions) (Phillips et al., 2004; Phillips, 2009). Previous studies on mitochondrial genomes have shown that RY-coding can improve the level of signal relative to compositional heterogeneity, by giving more weight to the slowest evolving sites in the mitochondrial genomes (Brinkmann and Philippe, 1999; Phillips and Penny, 2003; Delsuc et al., 2003; Phillips et al., 2004). This approach can therefore increase the ratio between historical and non-historical signal, reducing the weight of systematic biases. In order to control for this issue, we compiled and analyzed three additional datasets: (1) third codon position in protein-coding regions coded as RY [3RY], (2) positions with Cs and Ts replaced with Y [AGY], and (3) all nucleotides coded as purines (R) and pyrimidines (Y) [allRY]. For each dataset, relative support of internal nodes was assessed by bootstrap analyses using the bootstopping criteria autoMRE (Majority Rule Criterion) as implemented in RAxML 7.2.6 (Pattengale et al., 2009).

## 2.8. Taxon sampling effects on phylogenetic reconstruction

Incomplete taxon sampling has been demonstrated to be a major problem in phylogenetic reconstructions (Zwickl and Hillis, 2002; Hillis et al., 2003; Plazzi et al., 2010; Townsend and Leuenberger, 2011; Nabhan and Sarkar, 2012). Here, we explicitly test the importance of taxon representation in resolving the relationships between the major lineages within primates and colugos. We assembled four additional datasets, including different numbers of primate taxa in each: (1) only six primate taxa (one for each main lineage within primates: Lemuriformes, Chyromyiformes, Lorisiformes, Tarsiiformes, Platyrrhini, Catarrhini), (2) ten taxa as used by Schmitz et al. (2002a), (3) 14 taxa as used by Arnason et al. (2008), and (4) 26 taxa as in Matsui et al. (2009). We then compared the results obtained by these four datasets with our original dataset with 62 primate species. Phylogenetic analyses were run using RAxML 7.2.6 as fully described above.

## 2.9. Divergence date estimation and cross-validation analyses

Fossil constraints were carefully selected based on criteria for choosing appropriate calibration points reviewed elsewhere (Raum et al., 2005; Ho and Phillips, 2009; Parham et al., 2012). We employed seventeen fossil constraints identified from the literature and previous studies performed on primate and mammal phylogeny. Details about these fossil calibrations, including age, phylogenetic position, minimum and (putative) maximum bounds, and references are reported in Table 4. The detailed procedure of fossil selection, with paleontological details, and assignment to

alternative nodes for calibration of the molecular clock is fully reported in [Supplementary material B1](#).

We evaluated the congruence of these 17 minimum-age constraints using fossil cross-calibration (Near and Sanderson, 2004; Near et al., 2005; Hugall et al., 2007; Rutschmann et al., 2007). Fossil cross-validation involves the use of one fossil at a time to generate age estimates for the nodes to which the other fossils are assigned; then it compares the inferred age with the fossil age at that node. This technique assesses the consistency of date estimates generated from different putative calibration points (Near and Sanderson, 2004; Near et al., 2005). We calculated two different statistics to evaluate the congruence of each calibration point. First, following the method proposed by Near et al., we calculate the mean deviation ( $D$ ) between molecular (MA) and fossil age (FA) estimates for all nodes using a single fossil-dated node as a calibration point, and the SS values ( $SS_x = \sum_{i \neq x} D^2$ ) for a given fossil calibration node when it was used as the single calibration point. Second, we applied the statistics described by Hugall et al. (2007). We first calculate the mean deviation  $D$  as described above. Then, for each calibration point, we obtained the sum of the absolute values of differences between the estimated and proposed ages for each calibration node. This value ( $\sum |D|$ ) is an indication of how congruent a specific fossil calibration is with the remainder: the smaller the value the more consistent. For these analyses, we used the topology and branch lengths of the MT39 model tree obtained in our Bayesian analyses. This topology was strongly favored in the Bayes factor comparison analysis explained above (see [Table 3](#)).

For all the analyses described above, we used the rate-autocorrelated Bayesian relaxed molecular clock method implemented in the MULTIDIVTIME program package (Thorne and Kishino, 2002). First, maximum likelihood parameters for the F84 + G model of evolution were estimated with *baseml* from the PAML package

(Yang, 1997, 2007). Next, branch lengths and their variance-covariance matrix were calculated with *estbranches*. Finally divergence times were calculated with *multidivtime* using conservative priors and the fossil calibrations. Bayesian priors for *multidivtime* were chosen according to J. Thorne's recommendations in the *multidivtime* manual. Root-to-tip mean was set at 100.0 Ma, with a 100.0 Ma standard deviation. The evolutionary rate at the root was set at 0.009 substitutions per nucleotide site per Myr, with a standard deviation of 0.009. This was calculated as the median root-to-tip branch length divided by the root-to-tip mean prior. The *brownmean* and its standard deviation were set to 0.020, such that the *brownmean* multiplied by the root-to-tip mean prior (100 Ma) equals 2. After a burn-in period of 100,000 generations, MCMC chains were sampled every 100 generations until 10,000 samples were taken. To test the effect of the priors on the posteriors, several runs were performed with various prior choices. The results were extremely robust to changes in priors.

During the preliminary analyses fossil constraints were treated as point calibrations (minimum equal to maximum bounds). Based on the results of the preliminary analyses in which all 17 fossil calibrations were tested for congruence, a set of 16 constraints was used to obtain the final chronogram presented in the results. In this final analysis, for each fossil calibration we defined a minimum and a maximum constraint (if available) as reported in [Table 4](#).

### 3. Results

#### 3.1. Phylogeny

A comparison of the tree topologies that resulted from the analyses showed that varying partition scheme had little effect on the overall tree topology. For ML trees, between zero and six out of

**Table 4**  
Evolutionary rate calibration constraints (in millions of years).

Divergence	Min.	Max.	Fossil <sup>†</sup>	Reference	Age
<i>Primate fossils</i>					
1. <i>Homo</i>	5.0	10.0	<i>Ardipithecus</i> <i>Orrorin</i>	Haile-Selassie (2001) Senut et al. (2001)	5.2 6.0
2. <i>Gorilla</i>	10.0	NA	<i>Sahelanthropus</i> <i>Chororapithecus abyssinicus</i>	Vignaud et al. (2002) and Brunet et al. (2002) Suwa et al. (2007)	6.0–7.0 ≈10.0
3. <i>Pongo</i>	12.5	18.0	<i>Sivapithecus</i>	Kelley (2002)	≈12.5
4. Crown Catarrhini	21.0	33.9	<i>Morotopithecus</i> <i>Victoriapithecus</i>	Gebo et al. (1997) Pilbeam and Walker (1968) Benefit and McCrossin (2002)	>20.6 ≈19.0
5. <i>Theropithecus</i>	3.5	6.5	<i>Theropithecus</i>	Leakey (1993) Frost (2007)	≈3.5
6. <i>Macaca</i>	5.5	NA	<i>Macaca libyca</i>	Delson (1980) Jablonski (2002)	≈5.5
7. Colobinae	10	NA	<i>Microcolobus tugenensis</i>	Benefit and Pickford (1986) and Jablonski (2002)	9.0–11.0
8. <i>Aotus</i>	12.1	NA	<i>Aotus dindensis</i>	Setoguchi and Rosenberger (1987)	≈12.1
9. <i>Saimiri</i>	12.1	NA	<i>Neosaimiri</i>	Hartwig and Meldrum (2002)	≈12.1
10. Crown Anthropoidea	31.5	NA	Fayum catarrhines	Rasmussen (2002) and Seiffert et al. (2005)	31.5
11. Lorisioidea	36.9	NA	<i>Saharagalago</i> <i>Karanisia</i>	Seiffert et al. (2003) Seiffert (2007)	≈36.9
<i>Non-primate fossils</i>					
12. Cetartiodactyla	48.6	NA	<i>Himalayacetus</i> <i>Pakicetus</i>	Bajpai and Gingerich (1998) Gingerich and Russell (1981) Thewissen et al. (2001)	53.5 >48.6
13. Mysticeti-Odontoceti	33.5	53.5	<i>Llanocetus denticrenatus</i>	Mitchell (1989)	≈35
14. Feliniforma-Caniformia	43.0	63.8	<i>Tapocyon</i>	Benton and Donoghue (2007)	43.0–46.0
15. Bovidae	18.3	28.55	<i>Eotragus noyi</i>	Solounias et al. (1995) Benton and Donoghue (2007)	≈18.3
16. <i>Mus/Rattus</i>	7.3	16.0	<i>Progonomys hussaini</i>	Jacobs and Downs (1994) Benton and Donoghue (2007)	11.1–11.5
17. <i>Equus/Rhinoceros</i>	54.0	58.0	<i>Lambdotherium</i>	Rose (2006)	>50.3

<sup>†</sup> See [Supplementary material B1](#) for the detailed procedure of fossil selection, with paleontological details, and assignment to alternative nodes for calibration of the molecular clock.



168 possible conflicts were identified across partitions, representing a difference of only 0.00–3.57% (nodal distance: 0–0.474; split distance: 0–0.036). The differences were mainly represented by the position of two taxa: the genus *Cavia* within the rodents (either sister group of *Cricetulus*/*Mus/Rattus*) or sister taxon of the dormouse, *Glis glis*) and the position of *Cercopithecus diana* in relation to the other cercopithecids. Within the MB trees, only 2 possible conflicts were found across different partitioning scheme (1.07%; nodal distance: 0–0.331; split distance: 0–0.0119). The position of *C. diana* was the only taxon with conflicting placements among different partitions.

Bayes factor comparisons strongly supported the model with 39 partitions (MT39), consisting of two partitions for the rRNAs (12S and 16S), one for the 22 tRNAs, and 36 partitions for the three codon positions of each protein-coding gene. Partition scheme MT39 was strongly favored over MT6 (lnBF = 5531.809), MT6 was favored over MT5 (lnBF = 1287.011), MT5 strongly favored over MT15 (lnBF = 5068.416), MT15 favored over MT4 (lnBF = 996.090), and MT4 favored over MT0 (lnBF = 1440.723) (Table 3). The same results were obtained using maximum likelihood (MT39 < MT6 < MT5 < MT15 < MT4 < MT0). We therefore treat the MT39 ML and Bayesian trees as our “preferred tree”. Tree topology comparison analysis using TOPD-FMTS version 3.3 showed identical topologies between ML and Bayesian trees (0/168 nodes; Split Distance random [differeents/possibles]:  $0.997 \pm 0.007$  [ $167.46 \pm 1.195/168 \pm 0.000$ ]).

The Bayesian clade credibility values (PP) across the entire tree were high, with only two nodes below 0.95: the sister group relationship between *C. diana* and the clade including *Erythrocebus patas* and *C. mitis*/*C. lhoesti* (PP = 0.91), and the sister group relationship between *Erythrocebus patas* and *C. mitis*/*C. lhoesti* (PP = 0.80) (Fig. 1). In contrast to the Bayesian results, the maximum likelihood bootstrap (BP) values were quite low in several nodes across the tree (Fig. 1). Within primates several nodes were weakly supported in the ML analyses. For instance, within the platyrrhines, the sister relationship between *Ateles* and the clade including *Aotus*, *Saimiri*, *Cebus*, and *Saguinus* was below 70% bootstrap support. Also, several nodes within cercopithecids (particularly, the position of *E. patas* and *C. diana*, and the sister group relationship between *Nasalis* and *Pygathrix*) showed bootstrap support below 70% (Fig. 1). BP and PP values for all the nodes within primates are reported in Supplementary material (Table C1).

In order to evaluate the influence of RNA sequences on the topology and support values, we also ran RAxML and MrBayes using only the 12 protein-coding genes (11,022 bp). The topology of the reduced dataset did not differ from the one using all data. Similar to the results described above, BP values were generally lower than PP in Bayesian analyses (see Supplementary Fig. D1–2).

All the analyses supported the same topology and the monophyly of the order Primates (Fig. 1). Specifically, mitochondrial genome analyses supported the subdivision of primates into infraorders Strepsirrhini and Haplorhini, with tarsiers as sister group of Anthropoidea (BP > 70% and PP > 0.95). Moreover, the dermopterans (colugos, here represented by the genus *Galeopterus*) fell outside the primates (BP > 70% and PP > 0.95), and not the sister group of anthropoids as in many previous studies using mitochondrial DNA (Armason and Janke, 2002; Arnason et al., 2002, 2008; Schmitz et al., 2002a).

We evaluated the effect of compositional biases and the level of non-phylogenetic signal using three additional datasets (3RY, AGY, and allRY). All analyses strongly supported (BP > 70%) the monophyly of primates (with colugos outside primates), and two datasets (3RY and AGY) also strongly supported the sister group relationship between tarsiers and anthropoids. When all nucleotides were coded as RY (allRY), the monophyly of haplorhines

(tarsiers + anthropoids) was only weakly supported (BP = 57%) (Supplementary Fig. E1–3).

Relationships among the major lineages within primates were well supported by all the analyses, with lemuriformes and lorisooids within strepsirrhines, platyrrhines and catarrhines within anthropoids, and cercopithecoids (Cercopithecinae and Colobinae) and hominoids within catarrhines (Fig. 1). Two major discrepancies between our results and widely accepted relationships within primates are found in the phylogenetic relationships within the lorisooids and the papionins. In the former clade, the division between galagids and lorisooids was not supported by our mitogenomic data. The African lorisooids were found to be the sister group to the clade composed of Asian lorisooids and African galagids in all the analyses performed here. However, while Bayesian analyses strongly supported this topology (PP > 0.95), bootstrap values were relatively low across all the analyses (BP < 70%). Within the tribe Papionini, our results did not support the presence of an African clade (*Theropithecus/Papio/Lophocebus/Mandrillus/Cercocebus*). Instead, the group *Mandrillus/Cercocebus* clustered together with the genus *Macaca* (BP > 70% and PP > 0.95). Within these two primate groups (lorisooids and papionins), the same relationships were found when RY-coded datasets (3RY, AGY, and allRY) were analyzed.


### 3.2. Taxon sampling effects on phylogenetic reconstruction

In order to investigate the influence of taxon sampling in our dataset we analyzed four additional datasets including a subset of the primate taxa. Interestingly, all the datasets (6-taxa, 10-taxa, 14-taxa, and 26-taxa) showed no support for the monophyly of primates and the sister relationship between tarsiers and anthropoids. All reduced datasets supported instead the sister relationship between colugos and anthropoids and between strepsirrhines and tarsiers. However, the bootstrap values for both nodes were low (BP < 70%) across all the analyses: 32–60% for the clade colugos-anthropoids and 45–68% for the clade tarsiers-strepsirrhines (Supplementary Figs. F1–4).

### 3.3. Fossil congruence analyses

The cross-validation results (Table 5) reveal that calibration point 13 (C13: the split between Mysticeti and Odontoceti) is highly inconsistent with the other fossil calibrations. Specifically, when used as the only calibration point, C13 tended to overestimate all the other calibration points used in this study ( $D_x = 23.52$ ). Both statistics used in the study (SS and  $\Sigma D$ ) supported this conclusion (Table 5). All the other fossil calibrations used in this study provide plausible age estimates that fall within the range of the other calibrations. We thus used 16 calibration points listed in Table 4 (except C13) for all dating analyses.

Because the relative position of fossil calibrations in the tree might influence the precision of the dating estimates for the node of interest, we tested whether the absolute values of the deviation between the estimated and proposed ages for each calibration node were correlated with the branch lengths (nodal distance) between the calibration point used in each run and the remaining 16 nodes. We performed a Mantel test to check the correlation between the matrix of the deviations  $|D|$  (index of the estimation error) and the matrix of nodal distances in term of branch length (R Development Core Team, 2009). The Mantel test based on 10,000 replicates produced a positive correlation ( $p$ -value: 0.038) with  $r = 0.265$ . In order to evaluate the effect size of this correlation, we calculated the slope estimate from a linear regression. The analyses showed a positive relationship with  $\beta = 4.7$ .



**Fig. 1.** Single phylogenetic tree inferred from complete mitochondrial genome sequences based on maximum likelihood and Bayesian analyses. The tree topology remained constant across all analyses and branch lengths presented here follow the Bayesian analysis. The tree was rooted using *Loxodonta africana* and *Elephas maximus*. Maximum likelihood bootstrap values (BP) are shown with color coded inner circles (black: BP ≥ 90%, gray 70% ≤ BP < 90%; white: BP < 70%); Bayesian posterior probability value (PP) are indicated by the outer ring (black: PP ≥ 0.95, gray 0.75 ≤ PP < 0.95; white: PP < 0.75).

**3.4. Divergence dates**

Dating analyses were run with *multidivtime* using the MT39 partition strategy, as selected by the preliminary analyses de-

scribed above. Estimates for the major nodes within primates are reported in Table 6, along with 95% HPD intervals. The chronogram estimated the origin of crown primates at 74.1 Ma (95% HPD = 68.2–81.2; Fig. 2 and Table 6). Both crown strepsirrhines

**Table 5**

Cross-validation deviance of candidate calibration points. ( $D_x$ : average mean deviation between molecular and fossil age estimates for all available nodes based on the fossil calibration at node  $x$ ;  $SS$ : sum of the squared differences between the molecular and fossil age estimates at all other fossil-dated nodes;  $\Sigma D$ : sum of the absolute values of differences between the estimated and proposed ages for each calibration node).

CP	Divergence	Age	Node																	$D_x$	SS	$\Sigma D$
			C1	C2	C3	C4	C5	C6	C7	C8	C9	C10	C11	C12	C13	C14	C15	C16	C17			
C1	<i>Homo</i>	5.0	–	7.0	11.9	21.1	4.5	8.1	14.5	10.0	9.6	28.1	21.7	25.7	11.8	27.1	13.7	11.1	27.5	–5.60	2277.0	130.5
C2	<i>Gorilla</i>	10.0	7.2	–	17.0	30.4	6.5	11.6	20.9	14.4	13.8	40.6	31.4	37.1	17.0	39.2	19.8	16.0	39.7	–0.93	1099.6	110.94
C3	<i>Pongo</i>	12.5	5.4	7.5	–	21.9	4.8	8.5	15.2	10.4	10.0	29.0	22.3	26.2	11.9	27.7	13.9	11.3	28.1	–6.38	2181.2	127.77
C4	Crown Catarrhini	21.0	5.3	7.4	12.2	–	4.6	8.1	14.6	10.0	9.6	27.7	21.3	25.1	11.4	26.5	13.3	10.8	26.9	–5.21	2387.5	133.29
C5	<i>Theropithecus</i>	3.5	4.2	5.8	9.7	16.8	–	6.3	11.5	8.0	7.7	22.3	17.2	20.4	9.3	21.5	10.9	8.8	21.8	–9.75	3489.7	166.78
C6	<i>Macaca</i>	5.5	3.7	5.1	8.5	15.0	3.1	–	10.2	7.1	6.8	20.1	15.6	18.5	8.4	19.5	9.8	8.0	19.8	–11.56	4051.0	182.05
C7	Colobinae	10.0	3.6	5.0	8.4	14.5	3.1	5.6	–	6.9	6.6	19.2	14.8	17.5	7.9	18.5	9.3	7.5	18.7	–11.32	4336.2	188.3
C8	<i>Aotus</i>	12.1	6.6	9.2	15.2	26.3	5.7	10.2	18.3	–	11.7	35.0	27.1	32.2	14.8	33.9	17.2	13.9	34.4	–2.71	1373.5	110.84
C9	<i>Saimiri</i>	12.1	6.9	9.6	15.9	27.4	5.9	10.6	19.0	12.6	–	36.2	28.1	33.1	15.1	34.9	17.6	14.2	35.4	–1.88	1306.1	110.76
C10	Crown Anthroidea	31.5	6.0	8.4	13.9	23.9	5.2	9.3	16.7	11.3	10.9	–	24.3	28.9	13.2	30.4	15.3	12.5	30.8	–4.92	1772.1	117.83
C11	Lorisoidea	36.9	9.6	13.4	22.1	37.9	8.3	14.8	26.5	18.0	17.4	49.7	–	44.8	20.5	47.3	23.8	19.3	48.1	4.81	1599.1	139.06
C12	Cetartiodactyla	48.6	10.8	15.0	24.7	42.4	9.3	16.5	29.7	20.3	19.5	55.4	42.2	–	22.1	52.0	25.7	21.4	52.7	8.05	2393.3	168.76
C13	Mysticeti–Odontoceti	33.5	15.5	21.5	35.6	61.5	13.4	23.8	42.7	29.2	28.1	81.1	62.3	73.5	–	77.6	39.1	31.9	78.7	23.52	11026.8	383.95
C14	<i>Felis/Canis</i>	43.0	8.8	12.3	20.3	35.0	7.6	13.6	24.4	16.7	16.0	45.9	35.1	41.1	18.8	–	21.9	17.9	43.8	8.06	1313.0	125.72
C15	<i>Ovis/Bos</i>	18.3	7.3	10.2	16.8	29.0	6.3	11.2	20.2	13.8	13.3	38.2	29.3	34.4	15.7	36.3	–	14.8	36.8	–0.58	1251.1	113.95
C16	<i>Mus/Rattus</i>	7.3	4.0	5.6	9.3	16.0	3.5	6.2	11.1	7.6	7.3	21.0	16.1	19.1	8.8	20.2	10.2	–	20.5	–10.29	3836.8	174.83
C17	<i>Equus/Rhinoceros</i>	54.0	11.1	15.4	25.4	43.6	9.6	17.0	30.5	20.8	20.0	57.1	43.5	51.1	23.4	53.5	27.2	22.3	–	9.21	2694.5	180.89

and crown haplorhines are inferred to have originated in the Late Cretaceous, 66.3 Ma (95% HPD = 61.1–72.8) and 70.0 Ma (95% HPD = 64.3–76.7), respectively. The most recent common ancestor of the living platyrrhines was estimated to have lived 20.9 Ma (95% HPD = 17.9–24.4), while the origin of crown catarrhines occurred considerably earlier, approximately 32.1 Ma (95% HPD = 29.4–33.8). *Homo sapiens* and Neanderthals diverged at around 680,000 years ago (95% HPD = 490–926,000), while the date of the most recent common mitochondrial ancestor between the Denisovans, Neanderthals and modern humans was estimated to be approximately 1.39 Ma (95% HPD = 1.06–1.83). The estimates for all the nodes within primates are reported in the [Supplementary material \(Table C1\)](#).

#### 4. Discussion

In this study, we provided an assessment of primate phylogenetic relationships and divergence dates using complete mitochondrial genome data. We compiled one of the largest mitogenomic datasets for primates, and we applied a fossil cross-validation procedure to identify and select the most appropriate set of fossils to calibrate the molecular primate tree.

##### 4.1. Phylogenetic conclusion

All major lineages within the order Primates were included in our analyses and were inferred to be monophyletic. In contrast to previous mitogenomic studies, our analyses posit a monophyletic Primates with colugos as their sister group within Euarchontoglires. Previous mitogenomic studies hypothesized a sister relationship between anthropoid primates and colugos (Dermoptera), making the order Primates paraphyletic (e.g., [Arnason and Janke, 2002](#); [Arnason et al., 2002](#); [Schmitz et al., 2002a](#); [Kjer and Honeycutt, 2007](#)). [Arnason et al. \(2002\)](#) proposed the name Dermosimii for this assembly ([Arnason and Janke, 2002](#); [Arnason et al., 2002, 2008](#)). However, this group has been suggested to be an artifact of similar nucleotide composition of mitochondrial genomes rather than reflecting the true evolutionary history ([Schmitz et al., 2002a](#); [Schmitz and Zischler, 2003](#); [Raina et al., 2005](#); [Zischler, 2007](#)). In contrast to mitochondrial results, the monophyly of primates is in fact strongly supported not only by morphological studies ([Silcox et al., 2007](#)) but also by nuclear sequences ([Jameson](#)

[et al., 2011](#); [Meredith et al., 2011](#); [Perelman et al., 2011](#)) and transposable elements ([Schmitz et al., 2002a](#)). In this study we suggest that this artifact might also be a consequence of taxon sampling and the small internode distance between Dermoptera and Primates. Incomplete or biased taxon sampling is a major concern in phylogenetic studies and can provide misleading results in reconstructing phylogenetic relationships ([Zwickl and Hillis, 2002](#); [Hillis et al., 2003](#); [Plazzi et al., 2010](#); [Townsend and Leuenberger, 2011](#); [Nabhan and Sarkar, 2012](#)), and increased sampling of taxa has been shown to be an effective way to improve overall phylogenetic accuracy ([Zwickl and Hillis, 2002](#)). Here, we show that a large taxon sampling within primates can recover phylogenetic relationships consistent with morphological and nuclear data, and supports the monophyly of primates and the invalidity of the taxon Dermosimii. When only a subset of primate taxa was included in the analyses, the position of colugos and tarsiers became poorly supported and all the analyses recovered a sister relationship between colugos and anthropoids similar to previous studies (e.g., [Arnason and Janke, 2002](#); [Arnason et al., 2002](#); [Schmitz et al., 2002a](#); [Kjer and Honeycutt, 2007](#)), thus indicating that insufficient taxon sampling was likely the cause of the discrepancy found between mitochondrial DNA and other lines of evidence (e.g., [Schmitz et al., 2002a](#); [Silcox et al., 2007](#); [Jameson et al., 2011](#); [Meredith et al., 2011](#); [Perelman et al., 2011](#)).

The difficulty of recovering primate monophyly using mitochondrial sequences is somewhat confirmed by lower support in the ML bootstrap analyses, compared to Bayesian posterior probabilities. Theoretical and empirical studies showed that Bayesian posterior probability tends to be substantially higher than corresponding ML bootstrap frequencies. In particular, bootstrap analyses have been shown to be particularly sensitive to small numbers of characters and might underestimate confidence in tree topologies compared to Bayesian posterior probability in case of short internodes ([Alfaro, 2003](#); [Erixon et al., 2003](#); [Rothfels et al., 2012](#)). The low amount of phylogenetic signal in the mitochondrial genome for these nodes is demonstrated by the relatively short branch length of the internode between Dermoptera and Primates, and between Strepsirrhini and Haplorhini ([Fig. 1](#)).


Within primates, we found a well-supported division between strepsirrhines (including lorises, galagids, and Malagasy lemurs) and haplorhines (including tarsiers and anthropoids). Overall, support values across the entire primate tree were high with only a few exceptions. Within cercopithecoids, the sister group

**Table 6**

Comparison of divergence time estimates (posterior mean and 95% credible interval in millions of years) between this study and recent studies on primates.

Divergence (node)	This study	Arnason et al. (2008) <sup>c,d</sup>	Matsui et al. (2009) <sup>d</sup>	Fabre et al. (2009) <sup>e</sup>	Chatterjee et al. (2009) <sup>e</sup>	Perelman et al. (2011) <sup>f</sup>	Wilkinson et al. (2011) <sup>f</sup>	Jameson et al. (2011) <sup>f</sup>	Springer et al. (2012) <sup>e</sup>	Finstermeier et al. (2013) <sup>d</sup>
Crown Primates (61)	<b>74.1</b> <b>(68.2–81.2)</b>	80.0	76.0 (69.3–82.5)	78.8 (69.9–88.4)	63.7 (58.3–68.7)	87.2 (75.9–98.6)	84.5 (69.2–103.5)	72.6 (69.6–76.9)	67.8 (61.0–76.7)	66.2 (59.6–73.6)
Crown Strepsirrhini (60)	<b>66.3</b> <b>(61.1–72.8)</b>	70.0	64.5 (57.2–71.7)	67.1 (60.2–74.5)	51.6 (47.7–55.7)	68.7 (58.8–76.6)	49.8 (35.9–72.0)	52.4 (47.0–57.2)	54.2 (48.8–57.2)	56.9 (50.5–64.1)
Crown Lorisiformes (59)	<b>40.3<sup>a</sup></b> <b>(37.1–46.3)</b>	N/A	35.4 (28.5–43.1)	39.5 (38.0–41.8)	37.5 (36.9–38.7)	40.3 (35.2–45.6)	N/A	N/A	34.7 (27.9–38.2)	34.5 (30.2–39.0)
Lemuriformes–Chiromyiformes (55)	<b>59.6</b> <b>(54.3–65.9)</b>	64.0	55.3 (47.7–63.0)	59.6 (53.3–66.7)	46.2 (41.3–50.8)	58.6 (38.6–76.8)	N/A	N/A	50.0 (45.2–53.8)	47.1 (40.9–53.6)
Crown Lemuriformes (54)	<b>43.5</b> <b>(37.5–50.1)</b>	N/A	N/A	47.3 (37.9–50.8)	32.4 (28.6–33.6)	38.6 (26.4–50.0)	34.4 (23.5–49.1)	N/A	31.8 (26.7–36.6)	35.5 (30.4–40.4)
Crown Haplorhini (46)	<b>70.0</b> <b>(64.3–76.7)</b>	N/A	70.1 (63.2–76.7)	N/A	N/A	81.3 (69.5–95.8)	N/A	68.6 (65.6–72.7)	61.2 (57.6–69.6)	63.1 (56.0–69.7)
Crown Anthroidea (44)	<b>46.7<sup>a</sup></b> <b>(42.4–50.8)</b>	55.0	45.3 (39.4–51.3)	37.7 (33.3–42.7)	42.8 (40.1–45.0)	43.5 (38.6–48.4)	47.2 (38.9–56.5)	40.0 (37.3–43.1)	40.6 (33.6–49.5)	45.3 (40.7–50.1)
Crown Platyrrhini (43)	<b>20.9</b> <b>(17.9–24.4)</b>	N/A	N/A	14.5 (9.7–19.9)	26.6 (23.5–30.0)	24.8 (20.6–29.3)	25.1 (20.1–31.0)	N/A	23.3 (19.2–27.5)	22.0 (19.2–24.4)
Crown Catarrhini (38)	<b>32.1<sup>b</sup></b> <b>(29.4–33.8)</b>	39.0	30.5 (28.8–35.3)	23.9 (23.1–25.9)	29.3 (28.0–30.0)	31.6 (25.7–37.9)	31.0 (25.1–37.7)	25.4 (23.7–27.6)	25.1 (19.7–32.8)	31.9 (28.3–35.7)
Crown Cercopithecoidea (37)	<b>20.8<sup>a</sup></b> <b>(18.6–22.9)</b>	N/A	N/A	13.3 (11.6–14.7)	23.4 (21.9–24.9)	17.6 (13.9–21.5)	14.1 (11.0–17.7)	N/A	13.2 (8.9–18.3)	22.8 (20.0–25.6)
Crown Colobinae (36)	<b>14.1</b> <b>(12.0–16.3)</b>	N/A	N/A	8.7 (7.3–10.4)	18.4 (16.8–20.1)	12.3 (9.4–15.1)	N/A	N/A	8.9 (6.3–12.0)	15.5 (13.4–17.8)
Crown Cercopithecinae (29)	<b>14.1</b> <b>(12.2–15.8)</b>	N/A	N/A	9.1 (7.7–10.0)	18.6 (16.5–20.5)	11.5 (9.2–13.9)	10.3 (8.1–13.0)	N/A	8.4 (5.4–11.6)	14.9 (12.9–16.8)
Crown Hominoidea (12)	<b>22.3</b> <b>(20.5–23.9)</b>	N/A	19.9 (16.7–23.0)	18.6 (17.1–20.5)	21.5 (18.9–24.3)	20.3 (16.6–24.2)	19.2 (15.1–24.1)	N/A	17.4 (12.4–23.9)	20.3 (17.4–23.5)

<sup>a</sup> Lower bound defined as calibration point (see Table 4).<sup>b</sup> Lower and upper bound defined as calibration point (see Table 4).<sup>c</sup> Confidence intervals not provided.<sup>d</sup> The study used entire mitochondrial genomes.<sup>e</sup> The study used a supermatrix approach (mitochondrial and nuclear genes together).<sup>f</sup> The study used multiple nuclear loci (Perelman et al.: 54 loci ~35 kb – Wilkinson et al.: 2 genomic region ~83 kb – Jameson et al.: 1078 genes ~1.26 Mbp).



**Fig. 2.** Single chronogram with divergence date estimates from complete mitochondrial genome sequences. Only age estimates within primates are shown. Calibration points are depicted with filled circles; gray bars indicate constraint ranges. Mean node ages and 95% HPD intervals for major nodes within primates are presented in Table 6. Age estimates for all the nodes in the chronogram are reported in Supplementary material (Table C1).

relationship between *C. diana* and the clade including *Erythrocebus patas* and *C.mitis/C.lhoesti* and the sister group relationship between *Erythrocebus patas* and *C.mitis/C.lhoesti* were poorly supported in both Bayesian and ML analyses. Within the platyrrhines, most of the nodes showed low bootstrap support (but high Bayesian PP support), in agreement with previous studies that employed complete mitochondrial genomes (Hodgson et al., 2009; Chiou et al., 2011).

The phylogenetic position of the newly sequenced mitochondrial genomes was consistent with previous studies. The aye-aye

(*Daubentonia madagascariensis*) was inferred as the most basal divergence within lemurs, while *Mirza* (member of the family Cheirogaleidae) was recovered as sister group of sportive lemurs (*Lepilemur*) (Perelman et al., 2011; Springer et al., 2012). Within Cercopithecoidea, the phylogenetic positions of the five guenon genomes (*Allenopithecus nigroviridis*, *Cercopithecus diana*, *C. lhoesti*, *C. mitis*, and *Erythrocebus patas*) were identical to the ones suggested in a larger mitogenomic study for guenons (Guschanski et al., 2013), but in contrast with nuclear data (Perelman et al., 2011; but see Guschanski et al., 2013).

Despite the large taxon representation in this study, we could not overcome some well-known problems affecting mitochondrial-based reconstructions of primate phylogenetic relationships. For example, two well-supported taxonomic groups, the African papionins and the lorises, were not inferred to be monophyletic in our study. In the former group, our mitogenomic tree clusters *Mandrillus* and *Cercocebus* with *Macaca* instead of with the clade including *Papio*, *Theropithecus*, and *Lophocebus*. Within lorises, our mitogenomic tree places Asian lorises as the sister group to the African galagids rather than to the African lorisid *Perodicticus*. Both the lorises and the papionins are well supported by morphological data and nuclear sequences, and previous studies employing mitochondrial sequences have shown similar difficulty in recovering such clades (Papionini: Disotell et al., 1992; Harris, 2000; Finstermeier et al., 2013; Lorises: Roos et al., 2004; Masters et al., 2007; Matsui et al., 2009; Finstermeier et al., 2013). The mitochondrial genome represents a single locus and its phylogenetic history can differ from the species as a consequence of several events, including gene flow, hybridization, or incomplete lineage sorting (Maddison, 1997; Degnan and Rosenberg, 2006; Maddison and Knowles, 2006; Edwards, 2009). Although mitochondrial genomes are still a powerful resource to estimate phylogeny and phylogeography (Moore, 1995), conclusions based on a single locus should always be made with caution (Edwards, 2009; Ting and Sterner, 2013; Pozzi et al., 2014).

#### 4.2. Divergence dates

To our knowledge, this is the first study that explores the application of fossil cross-validation within primates. The primate fossil record is extremely fragmented and incomplete, and some studies estimate that only 4–7% of all primate species are known from the fossil record (Tavaré et al., 2002; Soligo and Martin, 2006; Martin et al., 2007). The degree of incompleteness of the fossil record is particularly extreme in some lineages, most notably within the Malagasy lemurs in which no fossils are known with the possible – but controversial – exception of *Bugtilemur mathesoni* in Pakistan (see Marivaux et al., 2001, 2006). As a consequence, most of the molecular studies to date employed a small set of calibration points, representing only a few nodes within the primate tree (e.g., Fabre et al., 2009; Matsui et al., 2009). More recently, Wilkinson et al. (2011) tried to better integrate the current knowledge of the fossil record into molecular date estimates. They developed a new methodology to better account for the rates of fossil preservation and discovery when calibrating a molecule-based tree. This study dated the origin for the crown primates at ~84 Ma, confirming a discrepancy between molecular estimates and the known fossil record.

In this study, we present a cross-validation analysis of the fossil record to select the most appropriate calibration points to be used with our dataset. Our approach consisted of two different steps: (1) we selected *a priori* putative fossils suitable to be used as calibration points for our dataset, following the criteria proposed by Parham et al. (2012); (2) we performed a cross-validation analyses aimed to detect possible outliers among our fossil selection. Since one of the main goals of this paper was to estimate the timing of primate origins we also included several calibration points within Boreoeutheria (Euarchontoglires and Laurasiatheria). Using only calibration points within primates might lead to problems in extrapolating dates, especially at the root of the tree (Raaijmakers et al., 2005). In this study we selected sixteen fossils as calibration points, including eleven within and five outside primates. The original cross-validation method proposed by Near et al. (2005) has been criticized by several authors because it could lead to several problems in selecting calibration points, including the possibility that the method will select an internally consistent set of fossils, exclud-

ing the most informative fossils because they are inconsistent with the majority of the remaining calibrations (Marshall, 2008; Ho and Phillips, 2009). Here, we used a conservative approach aimed to identify only possible outliers in the sets of fossils selected during the *a priori* phase. Although we acknowledge that the approach used here is not sufficient to address the complex challenges of calibration (Magallón, 2004; Near and Sanderson, 2004; Near et al., 2005; Huggall et al., 2007; Rutschmann et al., 2007; Pyron, 2010; Wilkinson et al., 2011), we believe that our approach allows researchers to better select fossils to be used in molecular phylogenetic analyses. For example, this approach might be beneficial in inferring the correct placement of fossils to specific nodes in a tree and excluding fossils that might be otherwise misused in calibrating molecular phylogenies. Moreover, new fossils can be tested against a good set of calibration points and alternative placements can be evaluated using the approach described here (see also Pyron, 2010 and Ward et al., 2010). This technique also allows researchers to include more fossils in their analyses, providing a better – and much needed – integration between molecular phylogenies and the paleontological record.

In order to identify how nodal distance can affect molecular estimates across the tree, we investigated the relationship between the deviation  $|D|$  and the nodal distance for each node selected as putative calibration point. Due to the incomplete nature of the fossil record, many areas of the primate tree have no available calibration point (e.g., the Malagasy lemurs). If the evolutionary distance from calibration points were a limiting factor in our ability to accurately estimate divergence times, then we would expect a strong correlation between nodal distance and the estimation error. The relationship that we found in our data set, although statistically significant, is particularly low. The Mantel test showed a weak positive correlation ( $r = 0.265$ ), suggesting that evolutionary distance from calibration points plays a very small role in the associated error. For instance, we found that the difference in estimated error of the crown Catarrhini divergence date between a near calibration point (*Homo/Pan*, branch length distance = 0.2588) and a more distant calibration point (Lorises/Galagidae, branch length = 0.7788) was only  $\pm 2.44$  million years. For this reason, our confidence in divergences distant from calibrations will be similar to those closer to calibration points. This result is particularly important for some taxonomic groups in which the fossil record is particularly scarce or absent. Therefore, despite the negative relationship between nodal distance and accuracy, modern molecular dating techniques can successfully provide reliable estimates even in areas where no calibration points are available.

The molecular chronogram obtained in this study estimated the crown-group origin of the order Primates in the Late Cretaceous (around 74 Ma), in agreement with most recent molecular studies (Arnason et al., 2008; Fabre et al., 2009; Matsui et al., 2009; Jameson et al., 2011; Perelman et al., 2011; Wilkinson et al., 2011). Nuclear estimates seem to provide slightly older estimates than the ones obtained by mitogenomic data (Perelman et al., 2011; Wilkinson et al., 2011), however all molecular data to date seem to strongly suggest an origin of primates before the K–Pg boundary. This result agrees with estimates obtained for other group of mammals, where the majority of recent molecular studies on mammalian diversification place the origin for most orders in the Cretaceous (Springer et al., 2003; Bininda-Emonds et al., 2007; Meredith et al., 2011).

A recent study on mitochondrial genomes including over 80 species of primates, recovered more recent origins for the crown group, dated around 66 Ma (59–73 Ma) (Finstermeier et al., 2013). This discordance with our study is particularly interesting given the large overlap in genome sequences used in both studies. Two main differences may have contributed to the discrepant conclusions between the two studies. First, Finstermeier et al. (2013)

only used calibration points within primates, making the age estimates at the root of the tree problematic due to extrapolation biases (see Raaum et al., 2005). The main differences in age estimates between Finstermeier et al.'s study and ours are limited to deeper nodes in the phylogeny. An additional source of discordance might be due to the use of controversial calibration points by Finstermeier et al. (2013). Out of the nine calibration points employed in that study, at least four were either secondary calibrations (i.e., molecular date estimates obtained from other molecular analyses) or based on misinterpretation of the fossil record (i.e., the use of stem fossils to calibrate a crown groups) (Graur and Martin 2004; Steiper and Young, 2008; Ho and Phillips, 2009; Pozzi et al., 2011). For instance, Finstermeier et al. (2013) calibrated the origins of crown catarrhines using the fossil *Saadanius hijazensis* (Zalmout et al., 2010). However, *Saadanius* is a stem catarrhine and cannot possibly inform the divergence between hominoids and catarrhines (see Pozzi et al., 2011).

Similar to Finstermeier et al. (2013), a recent study conducted by Steiper and Seiffert (2012) also suggested a more recent origin for crown primates, closer to the K–Pg boundary or possibly even in the Paleocene. In their study, the authors showed a negative correlation between three different life history variables (body size, absolute endocranial volume and relative endocranial volume) and molecular rates in primates. While almost all divergence date estimates for placental mammals based on molecular data are broadly consistent with our results, it is possible that there are life history factors that all these studies have failed to take into account (Bromham, 2009, 2011; Lanfear et al., 2010; Steiper and Seiffert, 2012).

Finally, O'Leary et al. (2013) recently integrated molecular and morphological evidence to support an explosive model of evolution for mammals, with most of the interordinal diversification dated just after the K–Pg boundary. Based on their analyses, the crown primates originated in the late Paleocene, around 55–56 Ma (O'Leary et al., 2013). The reconstruction proposed by O'Leary et al. (2013) is however based on phylogenetic analyses that fail to discriminate between homology and homoplasy and also implies extremely high acceleration in nucleotide substitution rate in early Paleocene mammals (Springer et al., 2013). One of the major shortcomings of O'Leary et al.'s study is the use of the earliest known fossil representative to date the origin of each group ('ghost lineage' analysis). As a consequence the origin of crown primates coincides with the earliest members of the group found in the fossil record at around 55–56 Ma (e.g., *Teilhardina brandti*) (O'Leary et al., 2013). However, the absence of evidence in the fossil record is not necessarily evidence of absence (Ho and Phillips, 2009) and the estimates obtained by O'Leary et al. (2013) can be considered only minimum divergence times based on ghost-lineage minimization rather than actual divergence times (Slater, 2013; Springer et al., 2013; Yoder, 2013; dos Reis et al., 2014).

The ancestor of the living catarrhines (apes and Old World monkeys) was estimated to have lived approximately 32 Ma. This result is in contrast to Zalmout et al. (2010) who suggested that the divergence for crown catarrhines dates after 29 Ma, based on the stem catarrhine *Saadanius hijazensis*. Inferring a date older than 29 Ma is not surprising because stem fossils do not provide any information about the divergence time of a crown group (Steiper and Young, 2008; Ho and Phillips, 2009); therefore, *Saadanius* does not inform the time of origin of crown catarrhines (Pozzi et al., 2011).

A recent study conducted by Langergraber et al. (2012) introduced a novel method to date divergence times within great apes without employing fossils as calibration points. This method relies on recent advances in the direct measurement of the mutation rate in humans (Awadalla et al., 2010; Roach et al., 2010; Sanders et al., 2012; Scally and Durbin, 2012) and in the generation times within

wild populations of chimpanzees and gorillas (Langergraber et al., 2012). Interestingly, the estimates obtained by Langergraber et al. are in line with the ones reported in this study. The only contrast with the above study is the age estimate for the divergence between bonobos (*Pan paniscus*) and common chimpanzee (*Pan troglodytes*) dated between 1.5 and 2.6 Ma by Langergraber et al. and 2.4–3.8 Ma in our study.

Finally, our estimates of 680 ka for Neanderthal and modern humans, and ~1.4 Ma for Denisovans and *Homo sapiens* are older – roughly 1.4x – than the ones obtained in the original paper by Krause et al. (2010) also using mitochondrial genomes. We identified the origin of this discrepancy in the different use of calibration points employed in the two studies. Krause et al. in fact assumed an average divergence of human and chimpanzee mtDNAs of 6 million years, while in our study we allowed this node to date between 5 and 10 Ma. We inferred this divergence to be at approximately 7.6 Ma (95% HPD = 6.7–8.8), which is substantially older than the age used by Krause et al. (2010). When the split between *Homo* and *Pan* was set at 6 Ma, we obtained age estimates that agree with Krause et al. (*H. sapiens*–Neanderthal: 530 ka [397–693]; *H. sapiens*–Denisovans: 1.08 Ma [0.87–1.35]). We contend that the mitochondrial divergences obtained in our study are more realistic than the one presented by Krause et al. (2010), because the common ancestor of modern humans and chimpanzees is likely to be older than 6 Ma, based on both molecular estimates (Jameson et al., 2011; Wilkinson et al., 2011; Springer et al., 2012) and the fossil record (Haile-Selassie, 2001; Senut et al., 2001; Brunet et al., 2002; Vignaud et al., 2002).

#### 4.3. Conclusion

Our study applied a fossil cross-validation procedure to identify a set of fossil calibrations to be used in dating a molecular phylogeny within primates. We used a two-step approach that includes an *a priori* selection of putative fossils and a *posteriori* cross-validation analysis to exclude possible outliers. The set of calibrations obtained by such an analysis can then be used to estimate the ages of the nodes of interest. We therefore recommend the use of these fossil calibrations for future studies aimed to estimate divergence times within primates. Also, our study showed a weak correlation between nodal distance and accuracy, suggesting that distant calibration points can be reliably used for dating areas of the trees where the fossil record is particularly scant.

The results obtained in this study agree with recent studies of primate divergence dates, supporting an origin of the order in the Late Cretaceous. This suggests that the early primate fossil record is likely poorly sampled. Although primate evolution may involve a slowdown in molecular rates, most molecular studies to date suggest that it is highly unlikely that primates – together with most mammalian orders – originated after the K–Pg boundary. Based on the age estimated obtained in our study we suggest a short-fuse model of primate origins, whereby relatively little time (<8 Myr) passed between the origin of the order Primates and the diversification of its major clades, Strepsirrhini, Tarsiiformes, and Anthropoidea.

#### Acknowledgments

We thank Terry Harrison, Eric Delson, Stephen Frost, and Christina Bergery for helpful comments and suggestions. We also want to thank three anonymous reviewers and the editor for their comments that strongly improve the quality of our study. This research was partly supported by National Institutes of Health Grants R01-GM060760 and R24-GM65580.

## Appendix A. Supplementary material

Supplementary data associated with this article can be found in the online version, at <http://dx.doi.org/10.1016/j.ympev.2014.02.023>.

## References

- Abascal, F., Zardoya, R., Telford, M.J., 2010. TranslatorX: multiple alignment of nucleotide sequences guided by amino acid translations. *Nucleic Acids Res.* 38, W7–W13.
- Alfaro, M.E., 2003. Bayes or bootstrap? A simulation study comparing the performance of Bayesian Markov chain Monte Carlo sampling and bootstrapping in assessing phylogenetic confidence. *Mol. Biol. Evol.* 20, 255–266.
- Anderson, S., De Bruijn, M.H.L., Coulson, A.R., Eperon, I.C., Sanger, F., Young, I.G., 1982. Complete sequence of bovine mitochondrial DNA conserved features of the mammalian mitochondrial genome. *J. Mol. Biol.* 156, 683–717.
- Arnason, U., Gullberg, A., 1993. Comparison between the complete mtDNA sequences of the blue and the fin whale, two species that can hybridize in nature. *J. Mol. Evol.* 37, 312–322.
- Arnason, U., Gullberg, A., Xu, X., 1996. A complete mitochondrial DNA molecule of the white-handed gibbon, *Hylobates lar*, and comparison among individual mitochondrial genes of all hominoid genera. *Hereditas* 124 (2), 185–189.
- Arnason, U., Janke, A., 2002. Mitogenomic analyses of eutherian relationships. *Cytogenet. Genome Res.* 96, 20–32.
- Arnason, U., Gullberg, A., Widegren, B., 1991. The complete nucleotide sequence of the mitochondrial DNA of the fin whale, *Balaenoptera physalus*. *J. Mol. Evol.* 33, 556–568.
- Arnason, U., Gullberg, A., Gretarsdottir, S., Ursing, B., 2000. The mitochondrial genome of the sperm whale and a new molecular reference for estimating eutherian divergence dates. *J. Mol. Evol.* 50, 569–578.
- Arnason, U., Adegoke, J.A., Bodin, K., Born, E.W., Esa, Y.B., Gullberg, A., Nilsson, M., Short, R.V., Xu, X., Janke, A., 2002. Mammalian mitogenomic relationships and the root of the eutherian tree. *Proc. Natl. Acad. Sci.* 99 (12), 8151–8156.
- Arnason, U., Gullberg, A., Janke, A., 2004. Mitogenomic analyses provide new insights into cetacean origin and evolution. *Gene* 333, 27–34.
- Arnason, U., Adegoke, J.A., Gullberg, A., Harley, E.H., Janke, A., Kullberg, M., 2008. Mitogenomic relationships of placental mammals and molecular estimates of their divergences. *Gene* 421, 37–51.
- Awadalla, P., Gauthier, J., Myers, R., Casals, F., Hamdan, F.F., Griffing, A.R., Côté, M., Henrion, E., Spiegelman, D., Tarabeux, J., Piton, A., Yang, Y., Boyko, A., Bustamante, C., Xiong, L., Rapoport, J.L., Addington, A.M., DeLisi, J.L.E., Krebs, M.-O., Joaber, R., Millet, B., Fombonne, E., Motton, L., Zilversmit, M., Keebler, J., Daoud, H., Marineau, C., Roy-Gagnon, M.-H., Dubé, M.-P., Eyre-Walker, A., Drapeau, P., Stone, E., Lafrenière, R.G., Rouleau, G., 2010. Direct measure of the de novo mutation rate in autism and schizophrenia cohorts. *Am. J. Hum. Genet.* 87, 316–324.
- Bajpai, S., Gingerich, P., 1998. A new Eocene archaeocete (Mammalia, Cetacea) from India and the time of origin of whales. *Proc. Natl. Acad. Sci. USA* 95, 15464–15468.
- Bayona-Bafaluy, M.P., 2003. Revisiting the mouse mitochondrial DNA sequence. *Nucleic Acids Res.* 31, 5349–5355.
- Beard, K.C., 1990. Gliding behaviour and palaeoecology of the alleged primate family Paromomyidae (Mammalia, Dermoptera). *Nature* 345, 340–341.
- Beard, K.C., 1993. Origin and evolution of gliding in Early Cenozoic Dermoptera (Mammalia, Primatomorpha). In: MacPhee, R.D.E. (Ed.), *Primates and their Relatives in Phylogenetic Perspective*. Plenum, pp. 63–90.
- Beard, K., Houde, P., 1989. An unusual assemblage of diminutive plesiadapiforms (Mammalia, ?Primates) from the early Eocene of the Clark's Fork Basin, Wyoming. *J. Vertebr. Paleontol.* 9, 37–41.
- Benefit, B.R., McCrossin, M.L., 2002. The Victoriapithecidae, Cercopithecoidea. In: Hartwig, W.C. (Ed.), *The Primate Fossil Record*. Cambridge University Press, Cambridge, UK, pp. 241–253.
- Benefit, B., Pickford, M., 1986. Miocene fossil cercopithecoids from Kenya. *Am. J. Phys. Anthropol.* 46, 441–464.
- Benton, M.J., Donoghue, P.C.J., 2007. Paleontological evidence to date the tree of life. *Mol. Biol. Evol.* 24, 26–53.
- Bininda-Emonds, O.R.P., Cardillo, M., Jones, K.E., MacPhee, R.D.E., Beck, R.M.D., Grenyer, R., Price, S.A., Vos, R.A., Gittleman, J.L., Purvis, A., 2007. The delayed rise of present-day mammals. *Nature* 446, 507–512.
- Bloch, J.I., Boyer, D.M., 2002. Grasping primate origins. *Science* 298, 1606–1610.
- Bloch, J.I., Silcox, M.T., 2001. New basicrania of Paleocene–Eocene Ignacius: re-evaluation of the Plesiadapiform–Dermopteran link. *Am. J. Phys. Anthropol.* 116, 184–198.
- Bloch, J.I., Silcox, M.T., Boyer, D.M., Sargis, E.J., 2007. New Paleocene skeletons and the relationship of plesiadapiforms to crown-clade primates. *Proc. Natl. Acad. Sci. USA* 104, 1159–1164.
- Briggs, A.W., Good, J.M., Green, R.E., Krause, J., Maricic, T., Stenzel, U., Lalueza-Fox, C., Rudan, P., Brajkovic, D., Kucan, Z., Gusic, I., Schmitz, R., Doronichev, V.B., Golovanova, L.V., De la Rasilla, M., Fortea, J., Rosas, A., Pääbo, S., 2009. Targeted retrieval and analysis of five Neandertal mtDNA genomes. *Science* 325, 318–321.
- Brinkmann, H., Philippe, H., 1999. Archaea sister group of Bacteria? Indications from tree reconstruction artifacts in ancient phylogenies. *Mol. Biol. Evol.* 16, 817–825.
- Bromham, L., 2009. Why do species vary in their rate of molecular evolution? *Biol. Lett.* 5, 401–404.
- Bromham, L., 2011. The genome as a life-history character: why rate of molecular evolution varies between mammal species. *Philos. Trans. Roy. Soc. Lond., Ser. B: Biol. Sci.* 366, 2503–2513.
- Brunet, M., Guy, F., Pilbeam, D., Mackaye, H.T., Likius, A., Ahounta, D., Beauvilain, A., Blondel, C., Bocherens, H., Boisserie, J.-R., De Bonis, L., Coppens, Y., Dejax, J., Denys, C., Düringer, P., Eisenmann, V., Fanone, G., Fronty, P., Geraads, D., Lehmann, T., Lihoreau, F., Louchart, A., Mahamat, A., Merceron, G., Mouchelin, G., Otero, O., Pelaez Campomanes, P., Ponce De Leon, M., Rage, J.-C., Sapanet, M., Schuster, M., Sudre, J., Tassy, P., Valentin, X., Vignaud, P., Viriot, P., Zazzo, A., Zollikofer, C., 2002. A new hominid from the Upper Miocene of Chad, Central Africa. *Nature* 418, 145–151.
- Burk, A., Douzery, E., 2002. The secondary structure of mammalian mitochondrial 16S rRNA molecules: refinements based on a comparative phylogenetic approach. *J. Mamm. Evol.* 9, 225–252.
- Cartmill, M., 1972. Arboreal adaptations and the origin of the order Primates. In: Tuttle, R.H. (Ed.), *The Functional and Evolutionary Biology of Primates*. Aldine-Atherton, pp. 97–122.
- Castresana, J., 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol. Biol. Evol.* 17, 540–552.
- Chan, Y.-C., Roos, C., Inoue-Murayama, M., Inoue, E., Shih, C.-C., Pei, K.J.-C., Vigilant, L., 2010. Mitochondrial genome sequences effectively reveal the phylogeny of *Hylobates gibbons*. *PLoS One* 5, e14419.
- Chatterjee, H.J., Ho, S.Y.W., Barnes, I., Groves, C., 2009. Estimating the phylogeny and divergence times of primates using a supermatrix approach. *BMC Evol. Biol.* 9, 259.
- Chiou, K.L., Pozzi, L., Lynch Alfaro, J.W., Di Fiore, A., 2011. Pleistocene diversification of living squirrel monkeys (*Saimiri* spp.) inferred from complete mitochondrial genome sequences. *Mol. Phylogenet. Evol.* 59, 736–745.
- Conroy, G.C., 1990. *Primate Evolution*. W.W. Norton and Co., New York, p. 492.
- Covert, H.H., 2002. The earliest fossil primates and the evolution of prosimians. In: Hartwig, W.C. (Ed.), *The Primate Fossil Record*. Cambridge University Press, Cambridge, UK, pp. 13–20.
- Cummings, M.P., Otto, S.P., Wakeley, J., 1995. Sampling properties of DNA sequence data in phylogenetic analysis. *Mol. Biol. Evol.* 12, 814–822.
- Degnan, J.H., Rosenberg, N.A., 2006. Discordance of species trees with their most likely gene trees. *PLoS Genet.* 2, e68.
- Delson, E., 1980. Fossil macaques, phyletic relationships and a scenario of deployment. In: Lindburg, D.G. (Ed.), *The Macaques: Studies in Ecology, Behavior and Evolution*. Van Nostrand, New York, pp. 10–30.
- Delsuc, F., Phillips, M.J., Penny, D., 2003. Comment on “Hexapod Origins: Monophyletic or Paraphyletic?”. *Science* 301, 1482.
- D’Erchia, A.M., Gissi, C., Pesole, G., Saccone, C., Arnason, U., 1996. The guinea-pig is not a rodent. *Nature* 381, 597–600.
- Disotell, T.R., Honeycutt, R.L., Ruvalo, M., 1992. Mitochondrial DNA Phylogeny of the Old-World monkeytribe papionini. *Mol. Biol. Evol.* 9, 1–13.
- Do, C.B., Woods, D.A., Batzoglou, S., 2006. CONTRAfold: RNA secondary structure prediction without physics-based models. *Bioinformatics* 22, e90–e98.
- Do, C.B., Foo, C., Batzoglou, S., 2008. A max-margin model for efficient simultaneous alignment and folding of RNA sequences. *Bioinformatics* 24, i68–i76.
- Dornburg, A., Beaulieu, J.M., Oliver, J.C., Near, T.J., 2011. Integrating fossil preservation biases in the selection of calibrations for molecular divergence time estimation. *Syst. Biol.* 60, 519–527.
- dos Reis, M., Donoghue, P.C.J., Yang, Z., 2014. Neither phylogenomic nor palaeontological data support a Palaeogene origin of placental mammals. *Biol. Lett.* 10, 20131003. <http://dx.doi.org/10.1098/rsbl.2013.1003>.
- Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32, 1792–1797.
- Edwards, S.V., 2009. Is a new and general theory of molecular systematics emerging? *Evolution* 63, 1–19.
- Erixon, P., Sennblad, B., Britton, T., Oxelman, B., 2003. Reliability of Bayesian posterior probabilities and bootstrap frequencies in phylogenetics. *Syst. Biol.* 52, 665–673.
- Fabre, P.-H., Rodrigues, A., Douzery, E.J.P., 2009. Patterns of macroevolution among Primates inferred from a supermatrix of mitochondrial and nuclear DNA. *Mol. Phylogenet. Evol.* 53, 808–825.
- Finstermeier, K., Zinner, D., Brameier, M., Meyer, M., Kreuz, E., Hofreiter, M., Roos, C., 2013. A mitogenomic phylogeny of living primates. *PLoS One* 8 (7), e69504.
- Flynn, S.M.C., Carr, S.M., 2007. Interspecies hybridization on DNA resequencing microarrays: efficiency of sequence recovery and accuracy of SNP detection in human, ape, and codfish mitochondrial DNA genomes sequenced on a human-specific MitoChip. *BMC Genomics* 8, 339.
- Friedlaender, J.S., Friedlaender, F.R., Hodgson, J.A., Stoltz, M., Koki, G., Horvat, G., Zhadanov, S., Schurr, T.G., Merriwether, D.A., 2007. Melanesian mtDNA complexity. *PLoS One* 2, e248.
- Frost, S., 2007. African Pliocene and Pleistocene cercopithecoid evolution and global climatic change. In: Bobe, R., Alemseged, Z. (Eds.), *Hominin Environments in the East African Pliocene: An assessment of the Faunal Evidence*. Springer, New York, pp. 51–76.
- Gebo, D.L., Maclatchy, L., Kityo, R., Deino, A., Kingston, J., Pilbeam, D., 1997. A hominoid genus from the early Miocene of Uganda. *Science* 276, 401–404.
- Gingerich, P.D., 1986. Early Eocene *Cantius torresi*—oldest primate of modern aspect from North America. *Nature* 319, 319–321.



- Gingerich, P.D., Russell, D.E., 1981. *Pakicetus inachus*, a new archaeocete (Mammalia, Cetacea) from the early–middle Eocene Kuldana Formation of Kohat (Pakistan). Contributions from the Museum of Paleontology, University of Michigan, vol. 25, No. 11, pp. 235–246.
- Gissi, C., Gullberg, A., Arnason, U., 1998. The complete mitochondrial DNA sequence of the rabbit, *Oryctolagus cuniculus*. Genomics 169, 161–169.
- Gissi, C., Reyes, A., Pesole, G., Saccone, C., 2000. Lineage-specific evolutionary rate in mammalian mtDNA. Mol. Biol. Evol. 17, 1022–1031.
- Gokey, N.G., Cao, Z., Pak, J.W., Lee, D., McKiernan, S.H., McKenzie, D., Weindruch, R., Aiken, J.M., 2004. Molecular analyses of mtDNA deletion mutations in microdissected skeletal muscle fibers from aged rhesus monkeys. Aging Cell 3, 319–326.
- Graur, D., Martin, W., 2004. Reading the entrails of chickens: molecular timescales of evolution and the illusion of precision. Trends Genet. 20, 80–86.
- Green, R.E., Malaspina, A., Krause, J., Briggs, A.W., Johnson, P.L.F., Uhler, C., Meyer, M., Good, J.M., Maricic, T., Stenzel, U., Prüfer, K., Siebauer, M., Burbano, H.A., Ronan, M., Rothberg, J.M., Egholm, M., Rudan, P., Brajković, D., Kučan, Z., Gusić, I., Wikström, M., Laakkonen, L., Kelso, J., Slatkin, M., Pääbo, S., 2008. A complete Neandertal mitochondrial genome sequence determined by high-throughput sequencing. Cell 134, 416–426.
- Guschanski, K., Krause, J., Sawyer, S., Valente, L.M., Bailey, S., Finstermeier, K., Sabin, R., Gillissen, E., Sonet, G., Nagy, Z.T., Lenglert, G., Mayer, F., Savolainen, V., 2013. Next-generation museumics disentangles one of the largest primate radiations. Syst. Biol. 62 (4), 539–554.
- Haile-Selassie, Y., 2001. Late Miocene hominids from the middle Awash, Ethiopia. Nature 412, 178–181.
- Harris, E.E., 2000. Molecular systematics of the old world monkey tribe papionini: analysis of the total available genetic sequences. J. Hum. Evol. 38, 235–256.
- Hartwig, W.C., Meldrum, D.J., 2002. Miocene platyrrhines of the northern Neotropics. In: Hartwig, W. (Ed.), The Primate Fossil Record. Cambridge University Press, Cambridge, UK, pp. 175–189.
- Hauf, J., Waddell, P., Chalwatzis, N., Joger, U., Zimmermann, F., 2000. The complete mitochondrial genome sequence of the African elephant (*Loxodonta africana*), phylogenetic relationships of Proboscidae to other mammals, and D-loop heteroplasmy. Zoology 102, 184–195.
- Hiendleder, S., Mainz, K., 1998. Mitochondrial DNA indicates that domestic sheep are derived from two different ancestral maternal sources: no evidence for contributions from urial and argali sheep. J. Hered. 89, 113–120.
- Hillis, D., Bull, J., 1993. An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analysis. Syst. Biol. 42, 182–192.
- Hillis, D.M., Pollock, D.D., McGuire, J.A., Zwickl, D.J., 2003. Is sparse taxon sampling a problem for phylogenetic inference? Syst. Biol. 52, 124–126.
- Ho, S.Y.W., Phillips, M.J., 2009. Accounting for calibration uncertainty in phylogenetic estimation of evolutionary divergence times. Syst. Biol. 58, 367–380.
- Hodgson, J.A., Sterner, K.N., Matthews, L.J., Burrell, I.A.S., Rachana, A.J., Raam, R.L., Stewart, C.-B., Disotell, T.R., 2009. Successive radiations, not stasis, in the South American primate fauna. Proc. Natl. Acad. Sci. USA 106, 5534–5539.
- Horai, S., Hayasaka, K., Kondo, R., Tsugane, K., Takahata, N., 1995. Recent African origin of modern humans revealed by complete sequences of hominoid mitochondrial DNAs. Proc. Natl. Acad. Sci. USA 92, 532–536.
- Huelsenbeck, J.P., Rannala, B., 2004. Frequentist properties of Bayesian posterior probabilities of phylogenetic trees under simple and complex substitution models. Syst. Biol. 53, 904–913.
- Huelsenbeck, J.P., Ronquist, F., 2001. MRBAYES: Bayesian inference of phylogenetic trees. Bioinformatics 17, 754–755.
- Hugall, A.F., Foster, R., Lee, M.S.Y., 2007. Calibration choice, rate smoothing, and the pattern of tetrapod diversification according to the long nuclear gene RAG-1. Syst. Biol. 56, 543–563.
- Inoue, J.G., Miya, M., Tsukamoto, K., Nishida, M., 2003. Evolution of the deep-sea gulper eel mitochondrial genomes: large-scale gene rearrangements originated within the eels. Mol. Biol. Evol. 20, 1917–1924.
- Inoue, J.G., Miya, M., Lam, K., Tay, B.-H., Danks, J.A., Bell, J., Walker, T.I., Venkatesh, B., 2010. Evolutionary origin and phylogeny of the modern holocephalans (Chondrichthyes: Chimaeriformes): a mitogenomic perspective. Mol. Biol. Evol. 27, 2576–2586.
- IUCN, 2013. IUCN Red List of Threatened Species. Version 2013.2. <www.iucnredlist.org>.
- Jablonski, N.G., 2002. Fossil Old World monkeys: the late Neogene radiation. In: Hartwig, W.C. (Ed.), The Primate Fossil Record. Cambridge University Press, Cambridge, UK, pp. 255–299.
- Jacobs, L.L., Downs, W.R., 1994. The evolution of murine rodents in Asia. In: Tomida, Y., Li, C.K., Setoguchi, T. (Eds.), Rodent and Lagomorph Families of Asian Origins and their Diversification. National Science Museum Monograph, Tokyo, pp. 149–156.
- Jameson, N.M., Hou, Z.-C., Sterner, K.N., Weckle, A., Goodman, M., Steiper, M.E., Wildman, D.E., 2011. Genomic data reject the hypothesis of a prosimian primate clade. J. Hum. Evol. 61, 295–305.
- Kass, R.E., Raftery, A.E., 1995. Bayes factors. J. Am. Stat. Assoc. 90, 773–795.
- Kay, R.F., Thorington, R.W., 1990. Eocene plesiadapiform shows affinities with flying lemur not primates. Nature 345, 342–344.
- Kay, R.F., Thewissen, J.G.M., Yoder, A.D., 1992. Cranial anatomy of *Ignacius graybullianus* and the affinities of the Plesiadapiformes. Am. J. Phys. Anthropol. 89, 477–498.
- Kelley, J., 2002. The hominoid radiation in Asia. In: Hartwig, W. (Ed.), The Primate Fossil Record. Cambridge University Press, Cambridge, pp. 369–384.
- Kim, K., Lee, S., Jeong, H., Ha, J., 1998. The complete nucleotide sequence of the domestic dog (*Canis familiaris*) mitochondrial genome. Mol. Phylogenet. Evol. 10, 210–220.
- Kjer, K.M., Honeycutt, R.L., 2007. Site specific rates of mitochondrial genomes and the phylogeny of eutheria. BMC Evol. Biol. 7, 8.
- Krause, J., Fu, Q., Good, J.M., Viola, B., Shunkov, M.V., Derevianko, A.P., Pääbo, S., 2010. The complete mitochondrial DNA genome of an unknown hominin from southern Siberia. Nature 464, 894–897.
- Lanfear, R., Welch, J.J., Bromham, L., 2010. Watching the clock: studying variation in rates of molecular evolution between species. Trends Ecol. Evol. 25, 495–503.
- Langergraber, K.E., Prüfer, K., Rowney, C., Boesch, C., Crockford, C., Fawcett, K., Inoue, E., Inoue-Muruyama, M., Mitani, J.C., Muller, M.N., Robbins, M.M., Schubert, G., Stoinski, T.S., Viola, B., Watts, D., Wittig, R.M., Wrangham, R.W., Zuberbühler, K., Pääbo, S., Vigilant, L., Prüfer, K., Zuberbühler, K., 2012. Generation times in wild chimpanzees and gorillas suggest earlier divergence times in great ape and human evolution. Proc. Natl. Acad. Sci. USA 109, 15716–15721.
- Leakey, M., 1993. Evolution of *Theropithecus* in the Turkana Basin. In: Jablonski, N.G. (Ed.), *Theropithecus: The Rise and Fall of a Primate Genus*. Cambridge University Press, Cambridge, UK, pp. 85–123.
- Lei, R., Shore, G.D., Brenneman, R.A., Engberg, S.E., Sitzmann, B.D., Bailey, C.A., Kimmel, L.M., Randriamampionona, R., Ranaivoarisoa, J.F., Louis, E.E., 2010. Complete sequence and gene organization of the mitochondrial genome for Hubbard's sportive lemur (*Lepilemur hubbardorum*). Gene 464, 44–49.
- Letsch, H.O., Kück, P., Stocsits, R.R., Misof, B., 2010. The impact of rRNA secondary structure consideration in alignment and tree reconstruction: simulated data and a case study on the phylogeny of hexapods. Mol. Biol. Evol. 27, 2507–2521.
- Li, D., Fan, L., Zeng, B., Yin, H., Zou, F., Wang, H., Meng, Y., King, E., Yue, B., 2009. The complete mitochondrial genome of *Macaca thibetana* and a novel nuclear mitochondrial pseudogene. Gene 429, 31–36.
- Lin, Y., Waddell, P., Penny, D., 2002. Pika and vole mitochondrial genomes increase support for both rodent monophyly and glires. Gene 294, 119–129.
- Lopez, J., Cevario, S., O'Brien, S., 1996. Complete nucleotide sequences of the domestic cat (*Felis catus*) mitochondrial genome and a transposed mtDNA tandem repeat (Numt) in the nuclear genome. Genomics 246, 229–246.
- Maddison, W.P., 1997. Gene trees in species trees. Syst. Biol. 46, 523.
- Maddison, W.P., Knowles, L.L., 2006. Inferring phylogeny despite incomplete lineage sorting. Syst. Biol. 55, 21.
- Madsen, O., Scally, M., Douady, C.J., Kao, D.J., DeBry, R.W., Adkins, R., Amrine, H.M., Stanhope, M.J., De Jong, W.W., Springer, M.S., 2001. Parallel adaptive radiations in two major clades of placental mammals. Nature 409, 610–614.
- Magallón, S., 2004. Dating lineages: molecular and paleontological approaches to the temporal framework of clades. Int. J. Plant Sci. 165.
- Marivaux, L., Welcomme, J.L., Antoine, P.O., Métais, G., Baloch, I.M., Benammi, M., Chaimanee, Y., Ducrocq, S., Jaeger, J.J., 2001. A fossil lemur from the Oligocene of Pakistan. Science (New York, NY) 294, 587–591.
- Marivaux, L., Chaimanee, Y., Tafforeau, P., Jaeger, J., 2006. New strepsirrhine primate from the late Eocene of Peninsular Thailand (Krabi Basin). Am. J. Phys. Anthropol. 130, 425–434.
- Marshall, C.R., 2008. A simple method for bracketing absolute divergence times on molecular phylogenies using multiple fossil calibration points. Am. Nat. 171, 726–742.
- Martin, R.D., 1968. Towards a new definition of Primates. Man New Series 3, 377–401.
- Martin, R., 1993. Primate origins: plugging the gaps. Nature 363, 223–234.
- Martin, R.D., Soligo, C., Tavaré, S., 2007. Primate origins: implications of a cretaceous ancestry. Folia Primatol. 78, 277–296.
- Mason, V.C., Li, G., Helgen, K.M., Murphy, W.J., 2011. Efficient cross-species capture hybridization and next-generation sequencing of mitochondrial genomes from noninvasively sampled museum specimens. Genome Res. 21, 1695–1704.
- Masters, J.C., Boniotto, M., Crovella, S., Roos, C., Pozzi, L., Delperio, M., 2007. Phylogenetic relationships among the Lorisioidea as indicated by craniodental morphology and mitochondrial sequence data. Am. J. Primatol. 69, 6–15.
- Matsudaira, K., Ishida, T., 2010. Phylogenetic relationships and divergence dates of the whole mitochondrial genome sequences among three gibbon genera. Mol. Phylogenet. Evol. 55, 454–459.
- Matsui, A., Rakotondraparany, F., Hasegawa, M., Horai, S., 2007. Determination of a complete lemur mitochondrial genome from feces. Mammal Study 32 (1), 7–16.
- Matsui, A., Rakotondraparany, F., Munechika, I., Hasegawa, M., Horai, S., 2009. Molecular phylogeny and evolution of prosimians based on complete sequences of mitochondrial DNAs. Gene 441, 53–66.
- Meredith, R.W., Janečka, J.E., Gatesy, J., Ryder, O., Fisher, C., Teeling, E.C., Goodbla, A., Eizirik, E., Simão, T.L.L., Stadler, T., Rabosky, D.L., Honeycutt, R.L., Flynn, J.J., Ingram, C.M., Steiner, C., Williams, T.L., Robinson, T.J., Burk-Herrick, A., Westerman, M., Ayoub, N., Springer, M.S., Murphy, W.J., 2011. Impacts of the cretaceous terrestrial revolution and KPg extinction on mammal diversification. Science (New York, NY) 334, 521–524.
- Miller, E.R., Gunnell, G.F., Martin, R.D., 2005. Deep time and the search for anthropoid origins. Am. J. Phys. Anthropol. 41 (Suppl.), 60–95.
- Mitchell, E.D., 1989. A new cetacean from the late Eocene La Meseta Formation, Seymour Island, Antarctic Peninsula. Can. J. Fish. Aquat. Sci. 46 (12), 2219–2235.
- Miya, M., Nishida, M., 2000. Use of mitogenomic information in teleostean molecular phylogenetics: a tree-based exploration under the maximum-parsimony optimality criterion. Mol. Phylogenet. Evol. 17, 437–455.
- Miya, M., Takeshima, H., Endo, H., Ishiguro, N.B., Inoue, J.G., Mukai, T., Satoh, T.P., Yamaguchi, M., Kawaguchi, A., Mabuchi, K., Shirai, S.M., Nishida, M., 2003. Major

- patterns of higher teleostean phylogenies: a new perspective based on 100 complete mitochondrial DNA sequences. *Mol. Phylogenet. Evol.* 26, 121–138.
- Moore, W., 1995. Inferring phylogenies from mtDNA variation: mitochondrial-gene trees versus nuclear-gene trees. *Evolution* 49, 718–726.
- Murphy, W.J., Eizirik, E., Johnson, W.E., Zhang, Y.P., Ryder, O.A., O'Brien, S.J., 2001. Molecular phylogenetics and the origins of placental mammals. *Nature* 409, 614–618.
- Nabhan, A.R., Sarkar, I.N., 2012. The impact of taxon sampling on phylogenetic inference: a review of two decades of controversy. *Brief. Bioinformatics* 13, 122–134.
- Near, T.J., Sanderson, M.J., 2004. Assessing the quality of molecular divergence time estimates by fossil calibrations and fossil-based model selection. *Philos. Trans. R. Soc. Lond. B: Biol. Sci.* 359, 1477–1483.
- Near, T.J., Meylan, P., Shaffer, H., 2005. Assessing concordance of fossil calibration points in molecular clock studies: an example using turtles. *Am. Nat.* 165.
- Newton, M., 1994. Approximate Bayesian Inference with the Weighted Likelihood Bootstrap. *J. Roy. Stat. Soc. B* 56, 3–48.
- Ni, X., Wang, Y., Hu, Y., Li, C., 2004. A euprimate skull from the early Eocene of China. *Nature* 427, 65–68.
- Nilsson, M.A., Gullberg, A., Spotorno, A.E., Arnason, U., Janke, A., 2003. Radiation of extant marsupials after the K/T boundary: evidence from complete mitochondrial genomes. *J. Mol. Evol.* 57 (Suppl. 1), S3–S12.
- Nylander, J., 2004. MrModeltest v2. Program Distributed by the author. Evolutionary Biology Centre, Uppsala University.
- Nylander, J., Ronquist, F., Huelsenbeck, J., Nieves-Aldrey, J., 2004. Bayesian phylogenetic analysis of combined data. *Syst. Biol.* 53, 47–67.
- Nylander, J., Wilgenbusch, J.C., Warren, D.L., Swofford, D.L., 2008. AWTY (are we there yet?): a system for graphical exploration of MCMC convergence in Bayesian phylogenetics. *Bioinformatics* 24, 581–583.
- O'Leary, M.A., Bloch, J.I., Flynn, J.J., Gaudin, T.J., Giallombardo, A., Giannini, N.P., Goldberg, S.L., Kraatz, B.P., Luo, Z.-X., Meng, J., Ni, X., Novacek, M.J., Perini, F.A., Randall, Z.S., Rougier, G.W., Sargis, E.J., Silcox, M.T., Simmons, N.B., Spaulding, M., Velazco, P.M., Weksler, M., Wible, J.R., Cirranello, A.L., 2013. The placental mammal ancestor and the Post-K–Pg radiation of placentals. *Science* 339, 662–667.
- Pacheco, M.A.M., Battistuzzi, F.F.U., Lentino, M., Aguilar, R., Kumar, S., Escalante, A.A., 2011. Evolution of modern birds revealed by Mitogenomics: timing the radiation and origin of major orders. *Mol. Biol. Evol.*
- Parham, J.F., Donoghue, P.C.J., Bell, C.J., Calway, T.D., Head, J.J., Holroyd, P.A., Inoue, J.G., Irmis, S.L., Joyce, W.G., Ksepka, D.T., Patané, J.S.L., Smith, N.D., Tarver, J.E., Van Tuinen, M., Yang, Z., Angielczyk, K.D., Greenwood, J.M., Hipsley, C.A., Jacobs, L., Makovicky, P.J., Müller, J., Smith, K.T., Theodor, J.M., Warnock, R.C.M., Benton, M.J., 2012. Best practices for justifying fossil calibrations. *Syst. Biol.* 61, 346–359.
- Partridge, M.A., Davidson, M.M., Hei, T.K., 2007. The complete nucleotide sequence of Chinese hamster (*Cricetus griseus*) mitochondrial DNA. *DNA Sequence: J. DNA Sequencing Mapp.* 18, 341–346.
- Pattengale, N.D., Alipour, M., Bininda-Emonds, O.R.P., Moret, B.M.E., Stamatakis, A., 2009. How many bootstrap replicates are necessary? In: *Research in Computational Molecular Biology*. Springer, Berlin Heidelberg, pp. 184–200.
- Pereira, S.L., Baker, A.J., 2006. A mitogenomic timescale for birds detects variable phylogenetic rates of molecular evolution and refutes the standard molecular clock. *Mol. Biol. Evol.* 23, 1731–1740.
- Perelman, P., Johnson, W.E., Roos, C., Seuánez, H.N., Horvath, J.E., Moreira, M.A.M., Kessing, B., Pontius, J., Roelke, M., Rumpel, Y., Schneider, M.P.C., Silva, A., O'Brien, S.J., Pecon-Slattery, J., 2011. A molecular phylogeny of living primates. *PLoS Genet.* 7, e1001342.
- Phillips, M.J., 2009. Branch-length estimation bias misleads molecular dating for a vertebrate mitochondrial phylogeny. *Gene* 441 (1), 132–140.
- Phillips, M.J., Penny, D., 2003. The root of the mammalian tree inferred from whole mitochondrial genomes. *Mol. Phylogenet. Evol.* 28, 171–185.
- Phillips, M.J., Delsuc, F., Penny, D., 2004. Genome-scale phylogeny and the detection of systematic biases. *Mol. Biol. Evol.* 21, 1455–1458.
- Pilbeam, D.R., Walker, A.C., 1968. Fossil monkeys from the Miocene of Napak, Northeast Uganda. *Nature* 220 (5168), 657–660.
- Piazza, F., Ferrucci, R.R., Passamonti, M., 2010. Phylogenetic representativeness: a new method for evaluating taxon sampling in evolutionary studies. *BMC Bioinformatics* 11, 209.
- Posada, D., Buckley, T.R., 2004. Model selection and model averaging in phylogenetics: advantages of akaike information criterion and bayesian approaches over likelihood ratio tests. *Syst. Biol.* 53, 793–808.
- Pozzi, L., Hodgson, J.A., Burrell, A.S., Disotell, T.R., 2011. The stem catarrhine *Saadanius* does not inform the timing of the origin of crown catarrhines. *J. Hum. Evol.* 61 (2), 209–210.
- Pozzi, L., Bergey, C.M., Burrell, A.S., 2014. The use (and misuse) of phylogenetic trees in comparative behavioral analyses. *Int. J. Primatol.* 35, 32–54.
- Puigbò, P., Garcia-Vallvé, S., McInerney, J.O., 2007. TOPD/FMTS: a new software to compare phylogenetic trees. *Bioinformatics* 23, 1556–1558.
- Pyron, R.A., 2010. A likelihood method for assessing molecular divergence time estimates and the placement of fossil calibrations. *Syst. Biol.* 59, 185–194.
- R Development Core Team, 2009. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. <<http://www.R-project.org>> (ISBN 3-900051-07-0).
- Raam, R.L., Sterner, K.N., Novello, C.M., Stewart, C.-B., Disotell, T.R., 2005. Catarrhine primate divergence dates estimated from complete mitochondrial genomes: concordance with fossil and nuclear DNA evidence. *J. Hum. Evol.* 48, 237–257.
- Raina, S.Z., Faith, J.J., Disotell, T.R., Seligmann, H., Stewart, C., Pollock, D.D., 2005. Evolution of base-substitution gradients in primate mitochondrial genomes. *Genome Res.* 15, 665–673.
- Rambaut, A., Drummond, A.J., 2007. Tracer v1.4. <<http://beast.bio.ed.ac.uk/Tracer>>.
- Rasmussen, D.T., 2002. Early catarrhines of the African Eocene and Oligocene. In: Hartwig, W.C. (Ed.), *The Primate Fossil Record*. Cambridge University Press, Cambridge, UK, pp. 203–220.
- Rasmussen, D.T., 2007. Fossil record of the primates from the Paleocene to the Oligocene. In: Henke, W., Tattersall, I. (Eds.), *Handbook of Paleoanthropology*. Springer, Berlin, pp. 889–920.
- Reyes, A., Pesole, G., Saccone, C., 1998. Complete mitochondrial DNA sequence of the fat dormouse, *Glis glis*: further evidence of rodent paraphyly. *Mol. Biol. Evol.* 15, 499–505.
- Reyes, A., Gissi, C., Pesole, G., 2000. Where do rodents fit? Evidence from the complete mitochondrial genome of *Sciurus vulgaris*. *Mol. Biol. Evol.* 17, 979–983.
- Roach, J.C., Glusman, G., Smit, A.F.A., Huff, C.D., Hubley, R., Shannon, P.T., Rowen, L., Pant, K.P., Goodman, N., Bamshad, M., Shendure, J., Drmanac, R., Jorde, L.B., Hood, L., Galas, D.J., 2010. Analysis of genetic inheritance in a family quartet by whole-genome sequencing. *Science (New York, NY)* 328, 636–639.
- Robinson, D., Foulds, L., 1981. Comparison of phylogenetic trees. *Math. Biosci.* 141, 131–141.
- Rogaev, E.I., Moliaka, Y.K., Malyarchuk, B.A., Kondrashov, F.A., Derenko, M.V., Chumakov, I., Grigorenko, A.P., 2006. Complete mitochondrial genome and phylogeny of Pleistocene mammoth *Mammuthus primigenius*. *PLoS Biol.* 4, e73.
- Ronquist, F., Huelsenbeck, J.P., 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19, 1572–1574.
- Roos, C., Schmitz, J., Zischler, H., 2004. Primate jumping genes elucidate strepsirrhine phylogeny. *Proc. Natl. Acad. Sci. USA* 101, 10650.
- Rose, K.D., 2006. *The Beginning of the Age of Mammals*. JHU Press, p. 428.
- Rothfels, C.J., Larsson, A., Kuo, L.-Y., Korall, P., Chiou, W.-L., Pryer, K.M., 2012. Overcoming deep roots, fast rates, and short internodes to resolve the ancient rapid radiation of eupolypod II ferns. *Syst. Biol.* 61, 490–509.
- Rowe, K.C., Singhal, S., Macmanes, M.D., Ayroles, J.F., Morelli, T.L., Rubidge, E.M., Bi, K., Moritz, C.C., 2011. Museum genomics: low-cost and high-accuracy genetic data from historical specimens. *Mol. Ecol. Resour.* 11, 1082–1092.
- Rutschmann, F., Eriksson, T., Salim, K.A., Conti, E., 2007. Assessing calibration uncertainty in molecular dating: the assignment of fossils to alternative calibration points. *Syst. Biol.* 56, 591–608.
- Sanders, S.J., Murtha, M.T., Gupta, A.R., Murdoch, J.D., Raubeson, M.J., Willsey, A.J., Ercan-Sencicek, A.G., DiLullo, N.M., Parikshak, N.N., Stein, J.L., Walker, M.F., Ober, G.T., Teran, N.A., Song, Y., El-Fishawy, P., Murtha, R.C., Choi, M., Overton, J.D., Bjornson, R.D., Carriero, N.J., Meyer, K.A., Bilguvar, K., Mane, S.M., Sestan, N., Lifton, R.P., Günel, M., Roeder, K., Geschwind, D.H., Devlin, B., State, M.W., 2012. De novo mutations revealed by whole-exome sequencing are strongly associated with autism. *Nature* 485, 237–241.
- Sally, A., Durbin, R., 2012. Revising the human mutation rate: implications for understanding human evolution. *Nat. Rev. Genet.* 13, 745–753.
- Schmitz, J., Zischler, H., 2003. A novel family of tRNA-derived SINEs in the colugo and two new retrotransposable markers separating dermopterans from primates. *Mol. Phylogenet. Evol.* 28, 341–349.
- Schmitz, J., Ohme, M., Suryobroto, B., Zischler, H., 2002a. The Colugo (*Cynocephalus variegatus*, Dermoptera): the primates' Gliding Sister? *Mol. Biol. Evol.* 19, 2308–2312.
- Schmitz, J., Ohme, M., Zischler, H., 2002b. The complete mitochondrial sequence of *Tarsius bancanus*: evidence for an extensive nucleotide compositional plasticity of primate mitochondrial DNA. *Mol. Biol. Evol.* 19, 544–553.
- Seiffert, E.R., 2007. Early evolution and biogeography of Lorisiform strepsirrhines. *Am. J. Primatol.* 69, 27–35.
- Seiffert, E.R., Simons, E.L., Attia, Y., 2003. Fossil evidence for an ancient divergence of lorises and galagos. *Nature* 422, 421–424.
- Seiffert, E.R., Simons, E.L., Clyde, W.C., Rossie, J.B., Attia, Y., Bown, T.M., Chatrath, P., Mathison, M.E., 2005. Basal anthropoids from Egypt and the antiquity of Africa's higher primate radiation. *Science (New York, NY)* 310, 300–304.
- Senut, B., Pickford, M., Gommery, D., Mein, P., Cheboi, K., Coppens, Y., 2001. First hominid from the Miocene (Lukeino Formation, Kenya). *C. R. Acad. Sci. - Ser. IIA - Earth Planet. Sci.* 332, 137–144.
- Setoguchi, T., Rosenberger, A.L., 1987. A fossil owl monkey from La Venta, Colombia. *Nature* 326, 692–694.
- Sigé, B., Jaeger, J.-J., Sudre, J., Vianey, Liaud, M., 1990. *Altiafasius koulchii* n. gen. et sp., primate omomyidid du Paléocène supérieur du Maroc, et les origines des euprimates. *Palaeontographica* 216, 31–56.
- Silcox, M.T., Sargis, E.J., Bloch, J.I., Boyer, D.M., 2007. Primate origins and supraordinal relationships: morphological evidence. In: Henke, W., Tattersall, I. (Eds.), *Handbook of Paleoanthropology*. Springer, New York, pp. 831–859.
- Slack, K.E., Delsuc, F., McLenachan, P., Arnason, U., Penny, D., 2007. Resolving the root of the avian mitogenomic tree by breaking up long branches. *Mol. Phylogenet. Evol.* 42, 1–13.
- Slater, G.J., 2013. Phylogenetic evidence for a shift in the mode of mammalian body size evolution at the Cretaceous–Palaeogene boundary. *Methods Ecol. Evol.* 4 (8), 734–744.
- Smith, T., Rose, K.D., Gingerich, P.D., 2006. Rapid Asia–Europe–North America geographic dispersal of earliest Eocene primate *Teilhardina* during the Paleocene–Eocene thermal maximum. *Proc. Natl. Acad. Sci. USA* 103, 11223–11227.
- Soligo, C., Martin, R.D., 2006. Adaptive origins of primates revisited. *J. Hum. Evol.* 50, 414–430.

- Solounias, N., Barry, J.C., Bernor, R.L., Lindsay, E.H., Raza, S.M., 1995. The oldest bovid from the Siwaliks, Pakistan. *J. Vertebr. Paleontol.* 15, 806–814.
- Song, S., Liu, L., Edwards, S.V., Wu, S., 2012. Resolving conflict in eutherian mammal phylogeny using phylogenomics and the multispecies coalescent model. *Proc. Natl. Acad. Sci. USA* 109, 14942–14947.
- Springer, M.S., Douzery, E., 1996. Secondary structure and patterns of evolution among mammalian mitochondrial 12S rRNA molecules. *J. Mol. Evol.* 43, 357–373.
- Springer, M.S., Murphy, W.J., Eizirik, E., O'Brien, S.J., 2003. Placental mammal diversification and the Cretaceous-Tertiary boundary. *Proc. Natl. Acad. Sci. USA* 100, 1056.
- Springer, M.S., Meredith, R.W., Gatesy, J., Emerling, C., Park, J., Rabosky, D.L., Stadler, T., Steiner, C., Ryder, O., Janečka, J.E., Fisher, C., Murphy, W.J., 2012. Macroevolutionary dynamics and historical biogeography of primate diversification inferred from a species supermatrix. *PLoS One* 7, e49521.
- Springer, M.S., Meredith, R.W., Teeling, E.C., Murphy, W.J., 2013. Technical comment on "the placental mammal ancestor and the post-K–Pg radiation of placentals". *Science* 341 (6146), 613.
- Stamatakis, A., 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22, 2688–2690.
- Stamatakis, A., Ludwig, T., Meier, H., 2005. RAxML-III: a fast program for maximum likelihood-based inference of large phylogenetic trees. *Bioinformatics* 21, 456–463.
- Stamatakis, A., Hoover, P., Rougemont, J., 2008. A rapid bootstrap algorithm for the RAxML Web servers. *Syst. Biol.* 57, 758–771.
- Steiper, M.E., Seiffert, E.R., 2012. Evidence for a convergent slowdown in primate molecular rates and its implications for the timing of early primate evolution. *Proc. Natl. Acad. Sci. USA* 109, 6006–6011.
- Steiper, M.E., Young, N.M., 2008. Timing primate evolution: Lessons from the discordance between molecular and paleontological estimates. *Evol. Anthropol.: Issues, News, Rev.* 17, 179–188.
- Stern, K.N., Raam, R.L., Zhang, Y.-P., Stewart, C.-B., Disotell, T.R., 2006. Mitochondrial data support an odd-nosed colobine clade. *Mol. Phylogenet. Evol.* 40, 1–7.
- Stocsits, R.R., Letsch, H., Hertel, J., Misof, B., Stadler, P.F., 2009. Accurate and efficient reconstruction of deep phylogenies from structured RNAs. *Nucleic Acids Res.* 37, 6184–6193.
- Suchard, M.A., Weiss, R.E., Sinsheimer, J.S., 2001. Bayesian selection of continuous-time Markov chain evolutionary models. *Mol. Biol. Evol.* 18, 1001–1013.
- Suwa, G., Kono, R.T., Katoh, S., Asfaw, B., Beyene, Y., 2007. A new species of great ape from the late Miocene epoch in Ethiopia. *Nature* 448, 921–924.
- Talavera, G., Castresana, J., 2007. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Syst. Biol.* 56, 564–577.
- Tavaré, S., Marshall, C.R., Will, O., Soligo, C., Martin, R.D., 2002. Using the fossil record to estimate the age of the last common ancestor of extant primates. *Nature* 416, 726–729.
- Thalmann, O., Hebler, J., Poinar, H.N., Pääbo, S., Vigilant, L., 2004. Unreliable mtDNA data due to nuclear insertions: a cautionary tale from analysis of humans and other great apes. *Mol. Ecol.* 13, 321–335.
- Thewissen, J.G.M., Williams, E.M., Hussain, S.T., 2001. Skeletons of terrestrial cetaceans and the relationship of whales to artiodactyls. *Nature* 413, 277–281.
- Thorne, J.L., Kishino, H., 2002. Divergence time and evolutionary rate estimation with multilocus data. *Syst. Biol.* 51, 689–702.
- Ting, N., Stern, K.N., 2013. Primate molecular phylogenetics in a genomic era. *Mol. Phylogenet. Evol.* 66, 565–568.
- Townsend, J.P., Leuenberger, C., 2011. Taxon sampling and the optimal rates of evolution for phylogenetic inference. *Syst. Biol.* 60, 358–365.
- Ursing, B.M., Arnason, U., 1998a. The complete mitochondrial DNA sequence of the pig (*Sus scrofa*). *J. Mol. Evol.* 47, 302–306.
- Ursing, B.M., Arnason, U., 1998b. Analyses of mitochondrial genomes strongly support a hippopotamus-whale clade. *Proceedings. Biol. Sci./Roy. Soc.* 265, 2251–2255.
- Ursing, B.M., Slack, K.E., Arnason, U., 2000. Subordinal artiodactyl relationships in light of phylogenetic analysis of 12 mitochondrial protein-coding genes. *Zool. Scr.* 29, 83–88.
- Vaidya, G., Lohman, D.J., Meier, R., 2011. SequenceMatrix: concatenation software for the fast assembly of multi-gene datasets with character set and codon information. *Cladistics* 27, 171–180.
- Vignaud, P., Douring, P., Mackaye, H.T., Likius, A., Blondel, C., Boisserie, J.-R., De Bonis, L., Eisenmann, V., Etienne, M.-E., Geraads, D., Guy, F., Lehmann, T., Lihoreau, F., Lopez-Martinez, N., Mourer-Chauviré, C., Otero, O., Rage, J.-C., Schuster, M., Viriot, L., Zazzo, A., Brunet, M., 2002. Geology and palaeontology of the Upper Miocene Toros-Menalla hominid locality, Chad. *Nature* 418, 152–155.
- Ward, P.S., Brady, S.G., Fisher, B.L., Schultz, T.R., 2010. Phylogeny and biogeography of dolichoderine ants: effects of data partitioning and relict taxa on historical inference. *Syst. Biol.* 59, 342–362.
- Wertheim, J.O., Worobey, M., 2007. A challenge to the ancient origin of SIVagm based on African green monkey mitochondrial genomes. *PLoS Pathog.* 3, e95.
- Wible, J., Covert, H.H., 1987. Primates: cladistic diagnosis and relationships. *J. Hum. Evol.* 16, 1–22.
- Wible, J.R., Rougier, G.W., Novacek, M.J., Asher, R.J., 2007. Cretaceous eutherians and Laurasian origin for placental mammals near the K/T boundary. *Nature* 447, 1003–1006.
- Wilcox, T.P., Zwickl, D.J., Heath, T.A., Hillis, D.M., 2002. Phylogenetic relationships of the dwarf boas and a comparison of Bayesian and bootstrap measures of phylogenetic support. *Mol. Phylogenet. Evol.* 25, 361–371.
- Wilkinson, R.D., Steiper, M.E., Soligo, C., Martin, R.D., Yang, Z., Tavaré, S., 2011. Dating primate divergences through an integrated analysis of palaeontological and molecular data. *Syst. Biol.* 60, 16–31.
- Williams, B.A., Kay, R.F., Kirk, E.C., 2010. New perspectives on anthropoid origins. *Proc. Natl. Acad. Sci. USA* 107, 4797–4804.
- Xu, X., Arnason, U., 1994. The complete mitochondrial DNA sequence of the horse, *Equus caballus*: extensive heteroplasmy of the control region. *Gene* 148, 357–362.
- Xu, X., Arnason, U., 1996. The mitochondrial DNA molecule of Sumatran orangutan and a molecular proposal for two (Bornean and Sumatran) species of orangutan. *J. Mol. Evol.* 43, 431–437.
- Xu, X., Janke, A., Arnason, U., 1996. The complete mitochondrial DNA sequence of the greater Indian rhinoceros, *Rhinoceros unicornis*, and the Phylogenetic relationship among Carnivora. *Mol. Biol. Evol.* 13, 1167–1173.
- Yang, Z., 1997. PAML: a program package for phylogenetic analysis by maximum likelihood. *Comput. Appl. Biosci.* CABIOS 13, 555–556.
- Yang, Z., 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* 24, 1586–1591.
- Yoder, A.D., 2013. Fossils versus clocks. *Science* 339, 656.
- Zalmout, I.S., Sanders, W.J., Maclatchy, L.M., Gunnell, G.F., Al-Mufarreh, Y.A., Ali, M.A., Nasser, A.-A.H., Al-Masari, A.M., Al-Sobhi, S.A., Nadhra, A.O., Matari, A.H., Wilson, J.A., Gingerich, P.D., 2010. New Oligocene primate from Saudi Arabia and the divergence of apes and Old World monkeys. *Nature* 466, 360–364.
- Zardoya, R., Malaga-Trillo, E., Veith, M., Meyer, A., 2003. Complete nucleotide sequence of the mitochondrial genome of a salamander, *Mertensiella luschani*. *Gene* 317, 17–27.
- Zhang, P., Wake, D.B., 2009. Higher-level salamander relationships and divergence dates inferred from complete mitochondrial genomes. *Mol. Phylogenet. Evol.* 53, 492–508.
- Zhang, P., Zhou, H., Liang, D., Liu, Y.-F., Chen, Y.-Q., Qu, L.-H., 2005. The complete mitochondrial genome of a tree frog, *Polypedates megacephalus* (Amphibia: Anura: Rhacophoridae), and a novel gene organization in living amphibians. *Gene* 346, 133–143.
- Zinner, D., Wertheimer, J., Liedigk, R., Groeneveld, L.F., Roos, C., 2013. Baboon phylogeny as inferred from complete mitochondrial genomes. *Am. J. Phys. Anthropol.* 150 (1), 133–140.
- Zischler, H., 2007. Molecular evidence on primate origins and evolution. In: Henke, W., Tattersall, I. (Eds.), *Handbook of Paleoanthropology*. Springer, New York, pp. 861–888.
- Zwickl, D.J., Hillis, D.M., 2002. Increased taxon sampling greatly reduces phylogenetic error. *Syst. Biol.* 51, 588–598.